

UNITED STATES PATENT APPLICATION

of

JOSE CIBELLI

ROBERT LANZA

and

JAMES M. ROBL

for

EMBRYONIC OR STEM-LIKE CELL LINES PRODUCED BY
CROSS SPECIES NUCLEAR TRANSPLANTATION AND METHODS
FOR ENHANCING EMBRYONIC DEVELOPMENT BY GENETIC
ALTERATION OF DONOR CELLS OR BY TISSUE CULTURE CONDITIONS

BURNS, DOANE, SWECKER & MATHIS, LLP
P.O. BOX 1404
ALEXANDRIA, VA 22313-1404

**EMBRYONIC OR STEM-LIKE CELL LINES PRODUCED BY
CROSS SPECIES NUCLEAR TRANSPLANTATION AND
METHODS FOR ENHANCING EMBRYONIC DEVELOPMENT
BY GENETIC ALTERATION OF DONOR CELLS OR
BY TISSUE CULTURE CONDITIONS**

CROSS-REFERENCE TO RELATES APPLICATIONS

This application claims priority under 35 U.S.C. §119 to PCT/US99/04608, filed on March 2, 1999. Also, this application is a continuation-in-part of U.S. Serial No. 09/395,368, filed September 14, 1999, which is a continuation-in-part of U.S. Serial No. 09/260,468, filed March 2, 1999, which is a continuation-in-part of U.S. Serial No. 09/032,945, filed March 2, 1998, which is in turn a continuation-in-part of U.S. Serial No. 08/699,040, filed on August 19, 1996. All of these applications are incorporated by reference in their entirety herein.

FIELD OF THE INVENTION

The present invention generally relates to the production of embryonic or stem-like cells by the transplantation of cell nuclei derived from animal or human cells into enucleated animal oocytes of a species different from the donor nuclei. The present invention more specifically relates to the production of primate or human embryonic or stem-like cells by transplantation of the nucleus of a primate or human cell into an enucleated animal oocyte, e.g., a primate or ungulate oocyte and in a preferred embodiment a bovine enucleated oocyte.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
1001
1002
1003
1004
1005
1006
1007
1008
1009
1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1020
1021
1022
1023
1024
1025
1026
1027
1028
1029
1030
1031
1032
1033
1034
1035
1036
1037
1038
1039
1040
1041
1042
1043
1044
1045
1046
1047
1048
1049
1050
1051
1052
1053
1054
1055
1056
1057
1058
1059
1060
1061
1062
1063
1064
1065
1066
1067
1068
1069
1070
1071
1072
1073
1074
1075
1076
1077
1078
1079
1080
1081
1082
1083
1084
1085
1086
1087
1088
1089
1090
1091
1092
1093
1094
1095
1096
1097
1098
1099
1100
1101
1102
1103
1104
1105
1106
1107
1108
1109
1110
1111
1112
1113
1114
1115
1116
1117
1118
1119
1120
1121
1122
1123
1124
1125
1126
1127
1128
1129
1130
1131
1132
1133
1134
1135
1136
1137
1138
1139
1140
1141
1142
1143
1144
1145
1146
1147
1148
1149
1150
1151
1152
1153
1154
1155
1156
1157
1158
1159
1160
1161
1162
1163
1164
1165
1166
1167
1168
1169
1170
1171
1172
1173
1174
1175
1176
1177
1178
1179
1180
1181
1182
1183
1184
1185
1186
1187
1188
1189
1190
1191
1192
1193
1194
1195
1196
1197
1198
1199
1200
1201
1202
1203
1204
1205
1206
1207
1208
1209
1210
1211
1212
1213
1214
1215
1216
1217
1218
1219
1220
1221
1222
1223
1224
1225
1226
1227
1228
1229
1230
1231
1232
1233
1234
1235
1236
1237
1238
1239
1240
1241
1242
1243
1244
1245
1246
1247
1248
1249
1250
1251
1252
1253
1254
1255
1256
1257
1258
1259
1260
1261
1262
1263
1264
1265
1266
1267
1268
1269
1270
1271
1272
1273
1274
1275
1276
1277
1278
1279
1280
1281
1282
1283
1284
1285
1286
1287
1288
1289
1290
1291
1292
1293
1294
1295
1296
1297
1298
1299
1300
1301
1302
1303
1304
1305
1306
1307
1308
1309
1310
1311
1312
1313
1314
1315
1316
1317
1318
1319
1320
1321
1322
1323
1324
1325
1326
1327
1328
1329
1330
1331
1332
1333
1334
1335
1336
1337
1338
1339
1340
1341
1342
1343
1344
1345
1346
1347
1348
1349
1350
1351
1352
1353
1354
1355
1356
1357
1358
1359
1360
1361
1362
1363
1364
1365
1366
1367
1368
1369
1370
1371
1372
1373
1374
1375
1376
1377
1378
1379
1380
1381
1382
1383
1384
1385
1386
1387
1388
1389
1390
1391
1392
1393
1394
1395
1396
1397
1398
1399
1400
1401
1402
1403
1404
1405
1406
1407
1408
1409
1410
1411
1412
1413
1414
1415
1416
1417
1418
1419
1420
1421
1422
1423
1424
1425
1426
1427
1428
1429
1430
1431
1432
1433
1434
1435
1436
1437
1438
1439
1440
1441
1442
1443
1444
1445
1446
1447
1448
1449
1450
1451
1452
1453
1454
1455
1456
1457
1458
1459
1460
1461
1462
1463
1464
1465
1466
1467
1468
1469
1470
1471
1472
1473
1474
1475
1476
1477
1478
1479
1480
1481
1482
1483
1484
1485
1486
1487
1488
1489
1490
1491
1492
1493
1494
1495
1496
1497
1498
1499
1500
1501
1502
1503
1504
1505
1506
1507
1508
1509
1510
1511
1512
1513
1514
1515
1516
1517
1518
1519
1520
1521
1522
1523
1524
1525
1526
1527
1528
1529
1530
1531
1532
1533
1534
1535
1536
1537
1538
1539
1540
1541
1542
1543
1544
1545
1546
1547
1548
1549
1550
1551
1552
1553
1554
1555
1556
1557
1558
1559
1560
1561
1562
1563
1564
1565
1566
1567
1568
1569
1570
1571
1572
1573
1574
1575
1576
1577
1578
1579
1580
1581
1582
1583
1584
1585
1586
1587
1588
1589
1590
1591
1592
1593
1594
1595
1596
1597
1598
1599
1600
1601
1602
1603
1604
1605
1606
1607
1608
1609
1610
1611
1612
1613
1614
1615
1616
1617
1618
1619
1620
1621
1622
1623
1624
1625
1626
1627
1628
1629
1630
1631
1632
1633
1634
1635
1636
1637
1638
1639
1640
1641
1642
1643
1644
1645
1646
1647
1648
1649
1650
1651
1652
1653
1654
1655
1656
1657
1658
1659
1660
1661
1662
1663
1664
1665
1666
1667
1668
1669
1670
1671
1672
1673
1674
1675
1676
1677
1678
1679
1680
1681
1682
1683
1684
1685
1686
1687
1688
1689
1690
1691
1692
1693
1694
1695
1696
1697
1698
1699
1700
1701
1702
1703
1704
1705
1706
1707
1708
1709
1710
1711
1712
1713
1714
1715
1716
1717
1718
1719
1720
1721
1722
1723
1724
1725
1726
1727
1728
1729
1730
1731
1732
1733
1734
1735
1736
1737
1738
1739
1740
1741
1742
1743
1744
1745
1746
1747
1748
1749
1750
1751
1752
1753
1754
1755
1756
1757
1758
1759
1760
1761
1762
1763
1764
1765
1766
1767
1768
1769
1770
1771
1772
1773
1774
1775
1776
1777
1778
1779
1780
1781
1782
1783
1784
1785
1786
1787
1788
1789
1790
1791
1792
1793
1794
1795
1796
1797
1798
1799
1800
1801
1802
1803
1804
1805
1806
1807
1808
1809
1810
1811
1812
1813
1814
1815
1816
1817
1818
1819
1820
1821
1822
1823
1824
1825
1826
1827
1828
1829
1830
1831
1832
1833
1834
1835
1836
1837
1838
1839
1840
1841
1842
1843
1844
1845
1846
1847
1848
1849
1850
1851
1852
1853
1854
1855
1856
1857
1858
1859
1860
1861
1862
1863
1864
1865
1866
1867
1868
1869
1870
1871
1872
1873
1874
1875
1876
1877
1878
1879
1880
1881
1882
1883
1884
1885
1886
1887
1888
1889
1890
1891
1892
1893
1894
1895
1896
1897
1898
1899
1900
1901
1902
1903
1904
1905
1906
1907
1908
1909
1910
1911
1912
1913
1914
1915
1916
1917
1918
1919
1920
1921
1922
1923
1924
1925
1926
1927
1928
1929
1930
1931
1932
1933
1934
1935
1936
1937
1938
1939
1940
1941
1942
1943
1944
1945
1946
1947
1948
1949
1950
1951
1952
1953
1954
1955
1956
1957
1958
1959
1960
1961
1962
1963
1964
1965
1966
1967
1968
1969
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
2024
2025
2026
2027
2028
2029
2030
2031
2032
2033
2034
2035
2036
2037
2038
2039
2040
2041
2042
2043
2044
2045
2046
2047
2048
2049
2050
2051
2052
2053
2054
2055
2056
2057
2058
2059
2060
2061
2062
2063
2064
2065
2066
2067
2068
2069
2070
2071
2072
2073
2074
2075
2076
2077
2078
2079
2080
2081
2082
2083
2084
2085
2086
2087
2088
2089
2090
2091
2092
2093
2094
2095
2096
2097
2098
2099
2100
2101
2102
2103
2104
2105
2106
2107
2108
2109
2110
2111
2112
2113
2114
2115
2116
2117
2118
2119
2120
2121
2122
2123
2124
2125
2126
2127
2128
2129
2130
2131
2132
2133
2134
2135
2136
2137
2138
2139
2140
2141
2142
2143
2144
2145
2146
2147
2148
2149
2150
2151
2152
2153
2154
2155
2156
2157
2158
2159
2160
2161
2162
2163
2164
2165
2166
2167
2168
2169
2170
2171
2172
2173
2174
2175
2176
2177
2178
2179
2180
2181
2182
2183
2184
2185
2186
2187
2188
2189
2190
2191
2192
2193
2194
2195
2196
2197
2198
2199
2200
2201
2202
2203
2204
2205
2206
2207
2208
2209
2210
2211
2212
2213
2214
2215
2216
2217
2218
2219
2220
2221
2222
2223
2224
2225
2226
2227
2228
2229
2230
2231
2232
2233
2234
2235
2236

In view of their ability to transfer their genome to the next generation, ES cells have potential utility for germline manipulation of livestock animals by using ES cells with or without a desired genetic modification. Moreover, in the case of livestock animals, e.g., ungulates, nuclei from like preimplantation livestock embryos support the development of enucleated oocytes to term (Smith et al., *Biol. Reprod.*, 40:1027-1035 (1989); and Keefer et al., *Biol. Reprod.*, 50:935-939 (1994)). This is in contrast to nuclei from mouse embryos which beyond the eight-cell stage after transfer reportedly do not support the development of enucleated oocytes (Cheong et al, *Biol. Reprod.*, 48:958 (1993)). Therefore, ES cells from livestock animals are highly desirable because they may provide a potential source of totipotent donor nuclei, genetically manipulated or otherwise, for nuclear transfer procedures.

Some research groups have reported the isolation of purportedly pluripotent embryonic cell lines. For example, Notarianni et al., *J. Reprod. Fert. Suppl.*, 43:255-260 (1991), report the establishment of purportedly stable, pluripotent cell lines from pig and sheep blastocysts which exhibit some morphological and growth characteristics similar to that of cells in primary cultures of inner cell masses isolated immunosurgically from sheep blastocysts. (*Id.*) Also, Notarianni et al., *J. Reprod. Fert. Suppl.*, 41:51-56 (1990) discloses maintenance and differentiation in culture of putative pluripotential embryonic cell lines from pig blastocysts. Further, Gerfen et al., *Anim. Biotech*, 6(1):1-14 (1995) disclose the isolation of embryonic cell lines from porcine blastocysts. These cells are stably maintained in mouse embryonic fibroblast feeder layers without the use of

conditioned medium. These cells reportedly differentiate into several different cell types during culture (Gerfen et al., *Id.*).

Further, Saito et al., *Roux's Arch. Dev. Biol.*, 201:134-141 (1992) report bovine embryonic stem cell-like cell lines cultured which survived passages for three, but were lost after the fourth passage. Still further, Handyside et al., *Roux's Arch. Dev. Biol.*, 196:185-190 (1987) disclose culturing of immunosurgically isolated inner cell masses of sheep embryos under conditions which allow for the isolation of mouse ES cell lines derived from mouse ICMs. Handyside et al. (1987) (*Id.*), report that under such conditions, the sheep ICMs attach, spread, and develop areas of both ES cell-like and endoderm-like cells, but that after prolonged culture only endoderm-like cells are evident. (*Id.*)

Recently, Cherny et al., *Theriogenology*, 41:175 (1994) reported purportedly pluripotent bovine primordial germ cell-derived cell lines maintained in long-term culture. These cells, after approximately seven days in culture, produced ES-like colonies which stain positive for alkaline phosphatase (AP), exhibited the ability to form embryoid bodies, and spontaneously differentiated into at least two different cell types. These cells also reportedly expressed mRNA for the transcription factors OCT4, OCT6 and HES1, a pattern of homeobox genes which is believed to be expressed by ES cells exclusively.

Also recently, Campbell et al., *Nature*, 380:64-68 (1996) reported the production of live lambs following nuclear transfer of cultured embryonic disc (ED) cells from day nine ovine embryos cultured under conditions which promote the isolation of ES cell lines

in the mouse. The authors concluded based on their results that ED cells from day nine ovine embryos are totipotent by nuclear transfer and that totipotency is maintained in culture.

Van Stekelenburg-Hamers et al., *Mol. Reprod. Dev.*, 40:444-454 (1995), reported the isolation and characterization of purportedly permanent cell lines from inner cell mass cells of bovine blastocysts. The authors isolated and cultured ICMs from 8 or 9 day bovine blastocysts under different conditions to determine which feeder cells and culture media are most efficient in supporting the attachment and outgrowth of bovine ICM cells. They concluded based on their results that the attachment and outgrowth of cultured ICM cells is enhanced by the use of STO (mouse fibroblast) feeder cells (instead of bovine uterus epithelial cells) and by the use of charcoal-stripped serum (rather than normal serum) to supplement the culture medium. Van Stekelenburg et al reported, however, that their cell lines resembled epithelial cells more than pluripotent ICM cells. (*Id.*)

Still further, Smith et al., WO 94/24274, published October 27, 1994, Evans et al, WO 90/03432, published April 5, 1990, and Wheeler et al, WO 94/26889, published November 24, 1994, report the isolation, selection and propagation of animal stem cells which purportedly may be used to obtain transgenic animals. Also, Evans et al., WO 90/03432, published on April 5, 1990, reported the derivation of purportedly pluripotent embryonic stem cells derived from porcine and bovine species which assertedly are useful for the production of transgenic animals. Further, Wheeler et al, WO 94/26884, published November 24, 1994, disclosed embryonic stem cells which are

assertedly useful for the manufacture of chimeric and transgenic ungulates. Thus, based on the foregoing, it is evident that many groups have attempted to produce ES cell lines, e.g., because of their potential application in the production of cloned or transgenic embryos and in nuclear transplantation.

5 The use of ungulate ICM cells for nuclear transplantation has also been reported. For example, Collas et al., *Mol. Reprod. Dev.*, 38:264-267 (1994) disclose nuclear transplantation of bovine ICMs by microinjection of the lysed donor cells into enucleated mature oocytes. The reference disclosed culturing of embryos *in vitro* for seven days to produce fifteen blastocysts which, upon transferral into bovine recipients, resulted in four pregnancies and two births. Also, Keefer et al., *Biol. Reprod.*, 50:935-939 (1994), disclose the use of bovine ICM cells as donor nuclei in nuclear transfer procedures, to produce blastocysts which, upon transplantation into bovine recipients, resulted in several live offspring. Further, Sims et al., *Proc. Natl. Acad. Sci., USA*, 90:6143-6147 (1993), disclosed the production of calves by transfer of nuclei from short-term *in vitro* cultured bovine ICM cells into enucleated mature oocytes.

Also, the production of live lambs following nuclear transfer of cultured embryonic disc cells has been reported (Campbell et al., *Nature*, 380:64-68 (1996)). Still further, the use of bovine pluripotent embryonic cells in nuclear transfer and the production of chimeric fetuses has also been reported (Stice et al., *Biol. Reprod.*, 54:100-110 (1996)); Collas et al, *Mol. Reprod. Dev.*, 38:264-267 (1994).

Further, there have been previous attempts to produce cross species NT units (Wolfe et al., *Theriogenology*, 33:350 (1990)). Specifically, bovine embryonic cells were fused with bison oocytes to produce some cross species NT units possibly having an inner cell mass. However, embryonic cells, not adult cells were used, as donor nuclei in the nuclear transfer procedure. The dogma has been that embryonic cells are more easily reprogrammed than adult cells. This dates back to earlier NT studies in the frog (review by DiBerardino, *Differentiation*, 17:17-30 (1980)). Also, this study involved very phylogenetically similar animals (cattle nuclei and bison oocytes). By contrast, previously when more diverse species were fused during NT (cattle nuclei into hamster oocytes), no inner cell mass structures were obtained. Further, it has never been previously reported that the inner cell mass cells from NT units could be used to form an ES cell-like colony that could be propagated.

Also, Collas et al (*Id.*), taught the use of granulosa cells (adult somatic cells) to produce bovine nuclear transfer embryos. However, unlike the present invention, these experiments did not involve cross-species nuclear transfer. Also, unlike the present invention ES-like cell colonies were not obtained.

Recently, U.S. Patent No. 5,843,780, issued to James A. Thomson on December 1, 1998, assigned to the Wisconsin Alumni Research Foundation, purports to disclose a purified preparation of primate embryonic stem cells that are (i) capable of proliferation in an *in vitro* culture for over one year; (ii) maintain a karyotype in which all chromosomes characteristic of the primate species are present and not noticeably altered

through prolonged culture; (iii) maintains the potential to differentiate into derivatives of endoderm, mesoderm and ectoderm tissues throughout culture; and (iv) will not differentiate when cultured on a fibroblast feeder layer. These cells were reportedly negative for the SSEA-1 marker, positive for the SEA-3 marker, positive for the SSEA-4 marker, express alkaline phosphatase activity, are pluripotent, and have karyotypes which include the presence of all the chromosomes characteristic of the primate species and in which none of the chromosomes are altered. Further, these cells are respectfully positive for the TRA-1-60, and TRA-1-81 markers. The cells purportedly differentiate into endoderm, mesoderm and ectoderm cells when injected into a SCID mouse. Also, purported embryonic stem cell lines derived from human or primate blastocysts are discussed in Thomson et al., *Science* 282:1145-1147 and *Proc. Natl. Acad. Sci., USA* 92:7844-7848 (1995).

Thus, Thomson disclose what purportedly are non-human primate and human embryonic or stem-like cells and methods for their production. However, there still exists a significant need for methods for producing human embryonic or stem-like cells that are autologous to an intended transplant recipient given their significant therapeutic and diagnostic potential.

In this regard, numerous human diseases have been identified which may be treated by cell transplantation. For example, Parkinson's disease is caused by degeneration of dopaminergic neurons in the substantia nigra. Standard treatment for Parkinson's involves administration of L-DOPA, which temporarily ameliorates the loss of dopamine,

but causes severe side effects and ultimately does not reverse the progress of the disease. A different approach to treating Parkinson's, which promises to have broad applicability to treatment of many brain diseases and central nervous system injury, involves transplantation of cells or tissues from fetal or neonatal animals into the adult brain. Fetal neurons from a variety of brain regions can be incorporated into the adult brain. Such grafts have been shown to alleviate experimentally induced behavioral deficits, including complex cognitive functions, in laboratory animals. Initial test results from human clinical trials have also been promising. However, supplies of human fetal cells or tissue obtained from miscarriages is very limited. Moreover, obtaining cells or tissues from aborted fetuses is highly controversial.

There is currently no available procedure for producing "fetal-like" cells from the patient. Further, allograft tissue is not readily available and both allograft and xenograft tissue are subject to graft rejection. Moreover, in some cases, it would be beneficial to make genetic modifications in cells or tissues before transplantation. However, many cells or tissues wherein such modification would be desirable do not divide well in culture and most types of genetic transformation require rapidly dividing cells.

There is therefore a clear need in the art for a supply of human embryonic or stem-like undifferentiated cells for use in transplants and cell and gene therapies.

OBJECTS OF THE INVENTION

It is an object of the invention to provide novel and improved methods for producing embryonic or stem-like cells.

It is a more specific object of the invention to provide a novel method for producing embryonic or stem-like cells which involves transplantation of the nucleus of a mammalian or human cell into an enucleated oocyte of a different species.

It is another specific object of the invention to provide a novel method for producing non-human primate or human embryonic or stem-like cells which involves transplantation of the nucleus of a non-human primate or human cell into an enucleated animal or human oocyte, e.g., an ungulate, human or primate enucleated oocyte.

It is another object of the invention to enhance the efficacy of cross-species nuclear transfer by introducing cytoplasm from an oocyte, blastomere or other embryonic cell that is compatible (same species) as the donor cell or nucleus.

It is a more specific object of the invention to enhance the efficacy of cross-species nuclear transfer by introducing cytoplasm from one or more immature oocytes which cytoplasm is compatible (same or closely related species) as the donor cell or nucleus. Such immature oocytes optionally may be matured *in vitro* and/or activated *in vitro* prior to collection and introduction of cytoplasm into the recipient enucleated oocyte.

It is another object of the invention to enhance the efficacy of cross-species nuclear transfer by incorporating mitochondrial DNA derived from the same or closely related species (preferably same donor) as the donor cell into the oocyte of a different species

that is used for nuclear transfer, before or after enucleation, or into the nuclear transfer unit (after the donor cell has been introduced).

It is still another object of the invention to enhance the efficacy of cross-species nuclear transfer by fusing an enucleated somatic cell (e.g., an enucleated human somatic cell) (karyoplast) with an activated or non-activated, enucleated or non-enucleated oocyte of a different species, e.g., bovine, or by fusion with an activated or unactivated cross-species NT unit which may be cleaved or uncleaved.

It is another object of the invention to provide a novel method for producing lineage-defective non-human primate or human embryonic or stem-like cells which involves transplantation of the nucleus of a non-human primate or human cell, e.g., a human adult cell into an enucleated non-human primate or human oocyte, wherein such cell has been genetically engineered to be incapable of differentiation into a specific cell lineage or has been modified such that the cells are "mortal", and thereby do not give rise to a viable offspring, e.g., by engineering expression of anti-sense or ribozyme telomerase gene.

It is still another object of the invention to enhance efficiency of nuclear transfer and specifically to enhance the development of preimplantation embryos produced by nuclear transfer by genetically engineering donor somatic cells used for nuclear transfer to provide for the expression of genes that enhance embryonic development, e.g., genes of the MHC I family, and in particular Ped genes such as Q7 and/or Q9.

It is another object of the invention to enhance the production of nuclear transfer embryos, e.g., cross-species nuclear transfer embryos, by the introduction of transgenes before or after nuclear transfer that provide for the expression of an antisense DNA encoding a cell death gene such as BAX, Apaf-1, or capsase, or a portion thereof, or demethylase.

It is yet another object of the invention to enhance the production of nuclear transfer embryos by IVP and more specifically nuclear transfer embryos by genetically altering the donor cell used for nuclear transfer such that it is resistant to apoptosis, e.g., by introduction of a DNA construct that provides for the expression of genes that inhibit apoptosis, e.g., Bcl-2 or Bcl-2 family members and/or by the expression of antisense ribozymes specific to genes that induce apoptosis during early embryonic development.

It is still another object of the invention to improve the efficacy of nuclear transfer by improved selection of donor cells of a specific cell cycle stage, e.g., G1 phase, by genetically engineering donor cells such that they express a DNA construct encoding a particular cyclin linked to a detectable marker, e.g., one that encodes a visualizable (e.g., fluorescent tag) marker protein.

It is also an object of the invention to enhance the development of *in vitro* produced embryos, by culturing such embryos in the presence of one or more protease inhibitors, preferably one or more capsase inhibitors, thereby inhibiting apoptosis.

It is another object of the invention to provide embryonic or stem-like cells produced by transplantation of nucleus of an animal or human cell into an enucleated oocyte of a different species.

5 It is a more specific object of the invention to provide primate or human embryonic or stem-like cells produced by transplantation of the nucleus of a primate or human cell into an enucleated animal oocyte, e.g., a human, primate or ungulate enucleated oocyte.

It is another object of the invention to use such embryonic or stem-like cells for therapy or diagnosis.

10 It is a specific object of the invention to use such primate or human embryonic or stem-like cells for treatment or diagnosis of any disease wherein cell, tissue or organ transplantation is therapeutically or diagnostically beneficial.

15 It is another specific object of the invention to use the embryonic or stem-like cells produced according to the invention for the production of differentiated cells, tissues or organs.

It is a more specific object of the invention to use the primate or human embryonic or stem-like cells produced according to the invention for the production of differentiated human cells, tissues or organs.

20 It is another specific object of the invention to use the embryonic or stem-like cells produced according to the invention for the production of genetically engineered embry-

onic or stem-like cells, which cells may be used to produce genetically engineered or transgenic differentiated human cells, tissues or organs, e.g., having use in gene therapies.

It is another specific object of the invention to use the embryonic or stem-like cells produced according to the invention *in vitro*, e.g. for study of cell differentiation and for assay purposes, e.g. for drug studies.

It is another object of the invention to provide improved methods of transplantation therapy, comprising the usage of isogenic or syngenic cells, tissues or organs produced from the embryonic or stem-like cells produced according to the invention. Such therapies include by way of example treatment of diseases and injuries including Parkinson's, Huntington's, Alzheimer's, ALS, spinal cord injuries, multiple sclerosis, muscular dystrophy, diabetes, liver diseases, heart disease, cartilage replacement, burns, vascular diseases, urinary tract diseases, as well as for the treatment of immune defects, bone marrow transplantation, cancer, among other diseases.

It is another object of the invention to use the transgenic or genetically engineered embryonic or stem-like cells produced according to the invention for gene therapy, in particular for the treatment and/or prevention of the diseases and injuries identified, *supra*.

It is another object of the invention to use the embryonic or stem-like cells produced according to the invention or transgenic or genetically engineered embryonic or stem-like cells produced according to the invention as nuclear donors for nuclear transplantation.

It is still another object of the invention to use genetically engineered ES cells produced according to the invention for the production of transgenic animals, e.g., non-human primates, rodents, ungulates, etc. Such transgenic animals can be used to produce, e.g., animal models for study of human diseases, or for the production of desired polypeptides, e.g., therapeutics or nutripharmaceuticals.

With the foregoing and other objects, advantages and features of the invention that will become hereinafter apparent, the nature of the invention may be more clearly understood by reference to the following detailed description of the preferred embodiments of the invention and to the appended claims.

BRIEFS DESCRIPTION OF THE FIGURES

Figure 1 is a photograph of a nuclear transfer (NT) unit produced by transfer of an adult human cell into an enucleated bovine oocyte.

Figures 2 to 5 are photographs of embryonic stem-like cells derived from a NT unit such as is depicted in Figure 1.

DETAILED DESCRIPTION OF THE INVENTION

The present invention provides a novel method for producing embryonic or stem-like cells, and more specifically non-human primate or human embryonic or stem-like cells by nuclear transfer or nuclear transplantation. In the subject application, nuclear transfer or nuclear transplantation or NT are used interchangeably.

As discussed *supra*, the isolation of actual embryonic or stem-like cells by nuclear transfer or nuclear transplantation has never been reported. Rather, previous reported isolation of ES-like cells has been from fertilized embryos. Also, successful nuclear transfer involving cells or DNA of genetically dissimilar species, or more specifically adult cells or DNA of one species (e.g., human) and oocytes of another non-related species has never been reported. Rather, while embryos produced by fusion of cells of closely related species, has been reported, e.g., bovine-goat and bovine-bison, they did not produce ES cells. (Wolfe et al, *Theriogenology*, 33(1):350 (1990).) Also, there has never been reported a method for producing primate or human ES cells derived from a non-fetal tissue source. Rather, the limited human fetal cells and tissues which are currently available must be obtained or derived from spontaneous abortion tissues and from aborted fetuses.

Also, prior to the present invention, no one obtained embryonic or stem-like cells by cross-species nuclear transplantation.

Quite unexpectedly, the present inventors discovered that human embryonic or stem-like cells and cell colonies may be obtained by transplantation of the nucleus of a human cell, e.g., an adult differentiated human cell, into an enucleated animal oocyte, which is used to produce nuclear transfer (NT) units, the cells of which upon culturing give rise to human embryonic or stem-like cells and cell colonies. This result is highly surprising because it is the first demonstration of effective cross-species nuclear transplantation involving the introduction of a differentiated donor cell or nucleus into an

enucleated oocyte of a genetically dissimilar species, e.g., the transplantation of cell nuclei from a differentiated animal or human cell, e.g., adult cell, into the enucleated egg of a different animal species, to produce nuclear transfer units containing cells which when cultured under appropriate conditions give rise to embryonic or stem-like cells and cell colonies.

Preferably, the recipient oocyte will be transplanted with a combination of (i) the donor cell or nucleus and (ii) at least one of compatible mitochondrial DNA or compatible cytoplasm. "Compatible" in the present application means that the mitochondrial DNA or cytoplasm is derived from a cell of the same species as the donor cell or nucleus, or a species that is very closely related to the donor cell or nucleus. For instance, if the donor cell or nucleus is a human cell or nucleus, the mitochondrial DNA or cytoplasm will be of human origin or that of a higher primate, e.g., chimpanzee, gorilla, or baboon. Similarly, if the donor cell or nucleus is a bovine cell or nucleus then compatible cytoplasm will be of a closely related ungulate, e.g., buffalo. Most preferably, the compatible mitochondrial DNA and/or cytoplasm will be derived from a cell of the same species as donor and most preferably will be autologous to the donor cell or nucleus.

In a preferred embodiment, the cytoplasm will be derived from immature oocytes or blastomeres, wherein such immature oocytes optionally may be matured *in vitro* and/or activated *in vitro* prior to collection of cytoplasm therefrom. Methods of *in vitro*

activating oocytes, including human oocytes, are described *infra*. Similar, methods of maturing oocytes *in vitro* have been reported in the literature.

Preferably, the NT units used to produce ES-like cells will be cultured to a size of at least 2 to 400 cells, preferably 4 to 128 cells, and most preferably to a size of at least about 50 cells.

In the present invention, embryonic or stem-like cells refer to cells produced according to the present invention. The present application refers to such cells as stem-like cells rather than stem cells because of the manner in which they are typically produced, i.e., by cross-species nuclear transfer. While these cells are expected to possess similar differentiation capacity as normal stem cells they may possess some insignificant differences because of the manner they are produced. For example, because these stem-like cells may possess the mitochondria of the oocytes used for nuclear transfer, and they may not behave or be morphogenetically identical to conventional embryonic stem cells.

The present discovery was made based on the observation that nuclear transplantation of the nucleus of an adult human cell, specifically a human epithelial cell obtained from the oral cavity of a human donor, when transferred into an enucleated bovine oocyte, resulted in the formation of nuclear transfer units, the cells of which upon culturing gave rise to human stem-like or embryonic cells and human embryonic or stem-like cell colonies. This result has recently been reproduced by transplantation of keratinocytes from an adult human into an enucleated bovine oocyte with the successful production of a blastocyst and ES cell line. Based thereon, it is hypothesized by the

present inventors that bovine oocytes and human oocytes, and likely mammals in general must undergo maturation processes during embryonic development which are sufficiently similar or conserved so as to permit the bovine oocyte to function as an effective substitute or surrogate for a human oocyte. Apparently, oocytes in general comprise factors, likely proteinaceous or nucleic acid in nature, that induce embryonic development under appropriate conditions, and these functions that are the same or very similar in different species. These factors may comprise material RNAs and/or telomerase.

Based on the fact that human cell nuclei can be effectively transplanted into bovine oocytes, it is reasonable to expect that human cells may be transplanted into oocytes of other non-related species, e.g., other ungulates as well as other animals. In particular, other ungulate oocytes should be suitable, e.g. pigs, sheep, horses, goats, etc. Also, oocytes from other sources should be suitable, e.g. oocytes derived from other primates, such as chimpanzees, macaques, baboons, gorilla, Rhesus monkey, amphibians, rodents, rabbits, guinea pigs, etc. Further, using similar methods, it should be possible to transfer human cells or cell nuclei into human oocytes and use the resultant blastocysts to produce human ES cells.

Therefore, in its broadest embodiment, the present invention involves the transplantation of an animal or human cell nucleus or animal or human cell into an oocyte (preferably enucleated) of an animal species different from the donor nuclei, by injection or fusion, and to optionally further inject into said recipient oocyte compatible mitochondrial DNA and/or cytoplasm, preferably of the same species as the donor cell

or nucleus, to produce an NT unit containing cells which may be used to obtain embryonic or stem-like cells and/or cell cultures. Enucleation (removal of endogenous oocyte nucleus) may be effected before or after nuclear transfer. For example, the invention may involve the transplantation of an ungulate cell nucleus or ungulate cell into an enucleated oocyte of another species, e.g., another ungulate or non-ungulate, by injection or fusion, which cells and/or nuclei are combined to produce NT units which are then cultured under conditions suitable to obtain multicellular NT units, preferably comprising at least about 2 to 400 cells, more preferably 4 to 128 cells, and most preferably at least about 50 cells. The cells of such NT units may be used to produce embryonic or stem-like cells or cell colonies upon culturing.

sub B' The preferred embodiment of the invention comprises the production of non-human primate or human embryonic or stem-like cells by transplantation of the nucleus of a donor human cell or a human cell into an enucleated human, primate, or non-primate animal oocyte, e.g., an ungulate oocyte, and in a preferred embodiment a bovine enucleated oocyte. Preferably, the enucleated oocyte will also be injected with human cytoplasm (e.g., from at least one immature or mature oocyte or blastomere), or fused with a karyoplast???? (enucleated human oocyte or blastomere, or that of a higher primate) and/or human mitochondrial DNA.

In general, the embryonic or stem-like cells will be produced by a nuclear transfer process comprising the following steps:

(i) obtaining desired human or animal cells to be used as a source of donor nuclei (which may be genetically altered);

(ii) obtaining oocytes from a suitable source, e.g. a mammal and most preferably a primate or an ungulate source, e.g. bovine,

5 (iii) enucleating said oocytes by removal of endogenous nucleus;

(iv) transferring the human or animal cell or nucleus into the enucleated oocyte of an animal species different than the donor cell or nuclei, e.g., by fusion or injection, wherein steps (iii) and (iv) may be effected in either order;

(v) culturing the resultant NT product or NT unit to produce multiple cell structures (embryoid structures having a discernible inner cell mass); and

(vi) culturing cells obtained from said embryos to obtain embryonic or stem-like cells and stem-like cell colonies.

As noted, the nuclear transfer process preferably will also include the introduction of compatible mitochondrial DNA and/or cytoplasm, i.e., derived from the same species or closely related species to the donor cell or nucleus. This may be effected before, 15 concurrent, or after introduction of the donor cell or nucleus into the recipient oocyte. Preferably, the compatible cytoplasm and/or mitochondrial DNA will be introduced proximate to introduction of the donor cell or nucleus into the recipient oocyte, i.e., within about twenty-four hours of such introduction, more preferably within about six 20 hours of such introduction, and most preferably within about two to four hours of such introduction.

Sources of compatible cytoplasm include in particular immature oocytes which can be obtained in large numbers from oocytes. For example, in the case of humans, oocytes can be obtained from ovaries of consenting donors, e.g., women undergoing hysterectomies.

5 Such immature oocytes optionally will be matured *in vitro* prior to collection of cytoplasm therefrom. Methods of effecting *in vitro* maturation of oocytes are described in U.S. Patent 5,945,577, incorporated by reference in its entirety herein.

10 Additionally, such immature oocytes can further optionally be activated *in vitro* prior to collection of cytoplasm therefrom. Methods for parthenogenetically activating oocytes, including human oocytes, are well known. Exemplary methods for effecting *in vitro* activation of oocytes are described in U.S. Serial No. 08/888,057, which is incorporated by reference in its entirety herein. Preferred methods for effecting *in vitro* activation include the use of DMAP and ionomycin, and the use of cycloheximide and ionomycin.

15 The amount of cytoplasm introduced into the recipient oocyte preferably will be an amount that increases its cell volume by at most five-fold, preferably no more than about two-fold, and most preferably no more than about one and a half-fold. This may be effected by injecting cytoplasm from compatible oocytes into the recipient oocyte or alternatively by fusing the recipient oocyte of compatible ooplasts (enucleated oocytes
20 of the same or very similar species as the recipient oocyte.) Optionally, all or part of the cytoplasm of the recipient oocyte may be removed to provide more cell volume for the

introduced compatible cytoplasm. With respect thereto, it has been suggested in the literature that maintaining appropriate nucleus/cytoplasm ratios and cell volume is importance to blastocyst formation and development. (Karyikova et al, *Reprod, Nutr. Devel.* 38(6):665-670)

5 While the present inventors do not wish to be bound by their belief, it is hypothesized that there are some constituents, e.g., nucleic acid sequences and/or proteins that are present in the cytoplasm of oocytes which facilitate embryogenesis while such factors are believed to be somewhat conserved across phylogenetically diverse species, it is theorized that the species differences are significant enough that they may impede the efficiency of cross-species nuclear transfer. Therefore, the present inventors hope to alleviate such efficacy problems by introducing compatible cytoplasm which should contain constituents of the same or very similar species that promote embryogenesis.

10 As noted above, cloning efficiency can also be improved by introducing compatible mitochondrial DNA, i.e., of the same or very similar species as the donor cell or nucleus. This can be effected before, concurrent, or after introduction of the donor cell or nucleus. Essentially, it is another object of the present invention to improve the efficacy of nuclear transfer, e.g., cross-species nuclear transfer by introducing mitochondrial DNA of the same species or highly similar species as the donor cell or nucleus into the recipient oocyte before or after nuclear transfer, before or after activation, and before or after fusion and cleavage. Preferably, if the donor cell is human, human

mitochondrial DNA will be derived from cells of the particular donor, e.g., liver cells and tissue.

Methods for isolating mitochondria are well known in the art. Mitochondria can be isolated from cells in tissue culture, or from tissue. The particular cells or tissue will depend upon the particular species of the donor cell. Examples of cells or tissues that may be used as sources of mitochondria include fibroblasts, epithelium, liver, lung, keratinocyte, stomach, heart, bladder, pancreas, esophageal, lymphocytes, monocytes, mononuclear cells, cumulus cells, uterine cells, placental cells, intestinal cells, hematopoietic cells, oocytes, and tissues containing such cells.

For example, mitochondria can be isolated from tissue culture cells and rat liver. It is anticipated that the same or similar procedures may be used to isolate mitochondria from other cells and tissues, i.e., human cells and tissues. As noted above, preferred source of mitochondria comprises human liver tissue because such cells contain a large number of mitochondria. Those skilled in the art will be able to modify the procedure as necessary, dependent upon the particular cell line or tissue. The isolated DNA can also be further purified, if desired, known methods, e.g., density gradient centrifugation.

The resultant isolated mitochondrial DNA will preferably be injected into the recipient oocyte. Preferably, the entire mitochondrial DNA isolated from one or several cells will be introduced to ensure that the recipient oocyte has a full complement of compatible mitochondrial DNA. Preferably, the mitochondrial DNA will be derived from an autologous source, i.e., cells or tissues obtained from the human subject who is the

source of the donor cell or nucleus, or a person genetically compatible therewith, e.g., close relative. Also, compatible mitochondria can be derived from immature oocytes and introduced together with the compatible cytoplasm.

The specific nuclear transfer or nuclear transplantation techniques that can be utilized in the present invention have been reported in the literature and are described in various of the references cited in the Background of the Invention. See, in particular, Campbell et al, *Theriogenology*, 43:181 (1995); Collas et al, *Mol. Report Dev.*, 38:264-267 (1994); Keefer et al, *Biol. Reprod.*, 50:935-939 (1994); Sims et al, *Proc. Natl. Acad. Sci., USA*, 90:6143-6147 (1993); WO 94/26884; WO 94/24274, and WO 90/03432, which are incorporated by reference in their entirety herein. Also, U.S. Patent Nos. 4,944,384 and 5,057,420 describe procedures for bovine nuclear transplantation. See, also Cibelli et al, *Science*, Vol. 280:1256-1258 (1998), U.S. Patent No. 5,945,577; WO 97/07669; WO 97/07668, and WO 96/07732, all of which are incorporated by reference in their entirety herein.

Human or animal cells, preferably mammalian cells, may be obtained and cultured by well known methods. Human and animal cells useful in the present invention include, by way of example, epithelial, neural cells, epidermal cells, keratinocytes, hematopoietic cells, melanocytes, chondrocytes, lymphocytes (B and T lymphocytes), other immune cells, erythrocytes, macrophages, melanocytes, monocytes, mononuclear cells, stem cells, fibroblasts, cardiac muscle cells, and other muscle cells, etc. Moreover, the human cells used for nuclear transfer may be obtained from different organs, e.g., skin, lung, pancreas,

liver, stomach, intestine, heart, reproductive organs, bladder, kidney, urethra and other urinary organs, etc. These are just examples of suitable donor cells.

Suitable donor cells, i.e., cells useful in the subject invention, may be obtained from any cell or organ of the body. This includes all somatic and germ cells. Preferably, the donor cells or nucleus will comprise an actively dividing, i.e., non-quiescent (G_1 , S or M cell cycle), cells as this has been reported to enhance cloning efficacy. Still more preferably, such donor cells will be in the G_1 cell cycle. However, quiescent donor cells (G_0) are also within the scope of the present invention. Quiescent cells may be obtained, e.g., appropriate culture conditions, by serum starvation.

The resultant blastocysts may be used to obtain embryonic stem cell lines according to the culturing methods reported by Thomson et al., *Science*, 282:1145-1147 (1998) and Thomson et al., *Proc. Natl. Acad. Sci., USA*, 92:7544-7848 (1995), incorporated by reference in their entirety herein.

Alternatively, the resultant blastocysts having a discernible inner cell mass can be cultured according to the method disclosed in U.S. Patent No. 5,905,042, incorporated by reference in its entirety herein. This method essentially comprises culturing of the entire inner cell mass or a portion thereof on a feeder layer, wherein contact with the feeder layer is maintained throughout culturing. This culturing method further comprises selective removal of a portion of the cultured inner cell mass as culturing proceeds, i.e., the outermost portion of the cultured inner cell mass that comprises cells possessing an ES-like appearance. This outermost portion of the cell colony is then introduced onto

another feeder layer, typically fetal fibroblasts. This method has been found to provide for the maintenance of non-differentiated (embryonic stem) cells in tissue culture, i.e., cells which exhibit an ES-like morphology which, upon removal from the feeder layer, differentiate into other cell types.

5 In the Example which follows, the cells used as donors for nuclear transfer were epithelial cells derived from the oral cavity of a human donor and adult human keratinocytes. However, as discussed, the disclosed method is applicable to other human cells or nuclei. Moreover, the cell nuclei may be obtained from both human somatic and germ cells.

10 It is also possible to arrest donor cells at mitosis before nuclear transfer, using a suitable technique known in the art. Methods for stopping the cell cycle at various stages have been thoroughly reviewed in U.S. Patent 5,262,409, which is herein incorporated by reference. In particular, while cycloheximide has been reported to have an inhibitory effect on mitosis (Bowen and Wilson (1955) *J. Heredity*, 45:3-9), it may also be
15 employed for improved activation of mature bovine follicular oocytes when combined with electric pulse treatment (Yang et al. (1992) *Biol. Reprod.*, 42 (Suppl. 1): 117).

20 Zygote gene activation is associated with hyperacetylation of Histone H4. Trichostatin-A has been shown to inhibit histone deacetylase in a reversible manner (Adenot et al, "Differential H4 acetylation of paternal and maternal chromatin precedes DNA replication and differential transcriptional activity in pronuclei of 1-cell mouse embryos." *Development* (Nov. 1997) 124(22): 4615-4625; Yoshida et al. "Trichostatin

A and trapoxin: novel chemical probes for the role of histone acetylation in chromatin structure and function." *Bioessays* (May, 1995) 17(5): 423-430), as have other compounds.

For instance, butyrate is also believed to cause hyper-acetylations of histones by inhibiting histone deacetylase. Generally, butyrate appears to modify gene expression and in almost all cases its addition to cell cultures appears to arrest cell growth. Use of butyrate to arrest cell growth is described in U.S. Patent No. 5,681,718, which is herein incorporated by reference. Thus, donor cells may be exposed to Trichostatin-A or another appropriate deacetylase inhibitor prior to fusion, or such a compound may be added to the culture media prior to genome activation.

Additionally, demethylation of DNA is thought to be a requirement for proper access of transcription factors to DNA regulatory sequences. Global demethylation of DNA from the eight-cell stage to the blastocyst stage in preimplantation embryos has previously been described (Stein et al., *Mol. Reprod. & Dev.*, 47(4): 421-429). Also, Jaenisch et al. (1997) have reported that 5-azacytidine can be used to reduce the level of DNA methylation in cells, potentially leading to increased access of transcription factors to DNA regulatory sequences. Accordingly, donor cells may be exposed to 5-azacytidine (5-Aza) previous to fusion, or 5-Aza may be added to the culture medium from the 8 cell stage to blastocyst. Alternatively, other known methods for effecting DNA demethylation may be used.

5 The oocytes used as the recipient for nuclear transfer or for recovery of compatible cytoplasm may be obtained from animals including mammals and amphibians. Suitable mammalian sources for oocytes include sheep, bovines, ovines, pigs, horses, rabbits, goats, guinea pigs, mice, hamsters, rats, primates, humans, etc. In the preferred
embodiments, the oocytes will be obtained from primates or ungulates, e.g., a bovine. In the case of human oocytes, a suitable source comprises ovaries obtained from consenting women, e.g., those undergoing hysterectomies.

10 Methods for isolation of oocytes are well known in the art. Essentially, this will comprise isolating oocytes from the ovaries or reproductive tract of a mammal or amphibian, e.g., a bovine. A readily available source of bovine oocytes is slaughterhouse materials.

15 For the successful use of techniques such as genetic engineering, nuclear transfer and cloning, oocytes preferably are matured *in vitro* before these cells can be used as recipient cells for nuclear transfer. This process generally requires collecting immature (prophase I) oocytes from animal ovaries, e.g., bovine ovaries obtained at a slaughterhouse and maturing the oocytes in a maturation medium prior to fertilization or enucleation until the oocyte attains the metaphase II stage, which in the case of bovine oocytes generally occurs about 18-24 hours post-aspiration. For purposes of the present
20 invention, this period of time is known as the "maturation period." As used herein for calculation of time periods, "aspiration" refers to aspiration of the immature oocyte from ovarian follicles.

Additionally, metaphase II stage oocytes, which have been matured *in vivo* have been successfully used in nuclear transfer techniques. For example, mature metaphase II oocytes can be collected surgically from either non-superovulated or superovulated cows or heifers 35 to 48 hours past the onset of estrus or past the injection of human chorionic gonadotropin (hCG) or similar hormone. Also, *in vivo* matured oocytes can be collected from other species, e.g., sheep, horse, goat pig, et al.

The stage of maturation of the oocyte at enucleation and nuclear transfer has been reported to be significant to the success of NT methods. (See e.g., Prather et al., *Differentiation*, 48, 1-8, 1991). In general, previous successful mammalian embryo cloning practices used metaphase II stage oocyte as the recipient oocyte because at this stage it is believed that the oocyte can be or is sufficiently "activated" to treat the introduced nucleus as it does a fertilizing sperm. In domestic animals, and especially cattle, the oocyte activation period generally ranges from about 16-52 hours, preferably about 28-42 hours post-aspiration.

For example, immature oocytes may be washed in HEPES buffered hamster embryo culture medium (HECM) as described in Seshagine et al., *Biol. Reprod.*, 40, 544-606, 1989, and then placed into drops of maturation medium consisting of 50 microliters of tissue culture medium (TCM) 199 containing 10% fetal calf serum which contains appropriate gonadotropins such as luteinizing hormone (LH) and follicle stimulating hormone (FSH), and estradiol under a layer of lightweight paraffin or silicon at 39°C.

After a fixed time maturation period, which typically will range from about 10 to 40 hours, and preferably about 16-18 hours, and which can occur *in vivo* or *in vitro*, the oocytes will preferably be enucleated. Prior to enucleation the oocytes will preferably be removed and placed in HECM containing 1 milligram per milliliter of hyaluronidase prior to removal of cumulus cells. This may be effected by repeated pipetting through very fine bore pipettes or by vortexing briefly. The stripped oocytes are then screened for polar bodies, and the selected metaphase II oocytes, as determined by the presence of polar bodies, are then used for nuclear transfer. Enucleation is typically effected at this time. However, enucleation may be effected before or after introduction of donor cell or nucleus because the donor nucleus is readily discernible from endogenous nucleus.

Enucleation may be effected by known methods, such as described in U.S. Patent No. 4,994,384 which is incorporated by reference herein. For example, metaphase II oocytes are either placed in HECM, optionally containing 7.5 micrograms per milliliter cytochalasin B, for immediate enucleation, or may be placed in a suitable medium, for example CR1aa, plus 10% estrus cow serum, and then enucleated later, preferably not more than 24 hours later, and more preferably 16-18 hours later.

Enucleation may be accomplished microsurgically using a micropipette to remove the polar body and the adjacent cytoplasm. The oocytes may then be screened to identify those of which have been successfully enucleated. This screening may be effected by staining the oocytes with 1 microgram per milliliter 33342 Hoechst dye in HECM, and then viewing the oocytes under ultraviolet irradiation for less than 10 seconds. The

oocytes that have been successfully enucleated can then be placed in a suitable culture medium.

In the present invention, the recipient oocytes will typically be enucleated at a time ranging from about 10 hours to about 40 hours after the initiation of *in vitro* maturation, more preferably from about 16 hours to about 24 hours after initiation of *in vitro* maturation, and most preferably about 16-18 hours after initiation of *in vitro* maturation. Enucleation may be effected before, simultaneous or after nuclear transfer. Also, enucleation may be effected before, after or simultaneous to activation.

A single animal or human cell or nucleus derived therefrom which is typically heterologous to the enucleated oocyte will then be transferred into the perivitelline space of the oocyte, typically enucleated, used to produce the NT unit. However, removal of endogenous nucleus may alternatively be effected after nuclear transfer. The animal or human cell or nucleus and the enucleated oocyte will be used to produce NT units according to methods known in the art. For example, the cells may be fused by electrofusion. Electrofusion is accomplished by providing a pulse of electricity that is sufficient to cause a transient break down of the plasma membrane. This breakdown of the plasma membrane is very short because the membrane reforms rapidly. Essentially, if two adjacent membranes are induced to break down, upon reformation the lipid bilayers intermingle and small channels will open between the two cells. Due to the thermodynamic instability of such a small opening, it enlarges until the two cells become one. Reference is made to U.S. Patent 4,997,384, by Prather et al., (incorporated by reference

in its entirety herein) for a further discussion of this process. A variety of electrofusion media can be used including e.g., sucrose, mannitol, sorbitol and phosphate buffered solution. Fusion can also be accomplished using Sendai virus as a fusogenic agent (Graham, *Wister Inot. Symp. Monogr.*, 9, 19, 1969).

5 Also, in some cases (e.g. with small donor nuclei) it may be preferable to inject the nucleus directly into the oocyte rather than using electroporation fusion. Such techniques are disclosed in Collas and Barnes, *Mol. Reprod. Dev.*, 38:264-267 (1994), and incorporated by reference in its entirety herein.

10 Preferably, the human or animal cell and oocyte are electrofused in a 500 μm chamber by application of an electrical pulse of 90-120V for about 15 μsec , about 24 hours after initiation of oocyte maturation. After fusion, the resultant fused NT units are preferably placed in a suitable medium until activation, e.g., one identified *infra*. Typically activation will be effected shortly thereafter, typically less than 24 hours later, and preferably about 4-9 hours later. However, it is also possible to activate the recipient
15 oocyte before or proximate (simultaneous) to nuclear transfer, which may or may not be enucleated. For example, activation may be effected from about twelve hours prior to nuclear transfer to about twenty-four hours after nuclear transfer. More typically, activation is effected simultaneous or shortly after nuclear transfer, e.g., about four to nine hours later.

20 The NT unit may be activated by known methods. Such methods include, e.g., culturing the NT unit at sub-physiological temperature, in essence by applying a cold, or

actually cool temperature shock to the NT unit. This may be most conveniently done by culturing the NT unit at room temperature, which is cold relative to the physiological temperature conditions to which embryos are normally exposed.

Alternatively, activation may be achieved by application of known activation agents. For example, penetration of oocytes by sperm during fertilization has been shown to activate prefusion oocytes to yield greater numbers of viable pregnancies and multiple genetically identical calves after nuclear transfer. Also, treatments such as electrical and chemical shock or cycloheximide treatment may also be used to activate NT embryos after fusion. Suitable oocyte activation methods are the subject of U.S. Patent No. 5,496,720, to Susko-Parrish et al., which is herein incorporated by reference.

For example, oocyte activation may be effected by simultaneously or sequentially:

- (i) increasing levels of divalent cations in the oocyte, and
- (ii) reducing phosphorylation of cellular proteins in the oocyte.

This will generally be effected by introducing divalent cations into the oocyte cytoplasm, e.g., magnesium, strontium, barium or calcium, e.g., in the form of an ionophore. Other methods of increasing divalent cation levels include the use of electric shock, treatment with ethanol and treatment with caged chelators.

Phosphorylation may be reduced by known methods, e.g., by the addition of kinase inhibitors, e.g., serine-threonine kinase inhibitors, such as 6-dimethylaminopurine, staurosporine, 2-aminopurine, and sphingosine.

Alternatively, phosphorylation of cellular proteins may be inhibited by introduction of a phosphatase into the oocyte, e.g., phosphatase 2A and phosphatase 2B.

Specific examples of activation methods are listed below.

1. Activation by Ionomycin and DMAP

- 1- Place oocytes in Ionomycin ($5\ \mu\text{M}$) with 2 mM of DMAP for 4 minutes;
- 2- Move the oocytes into culture media with 2 mM of DMAP for 4 hours;
- 3- Rinse four times and place in culture.

2. Activation by Ionomycin DMAP and Roscovitin

- 1- Place oocytes in Ionomycin ($5\ \mu\text{M}$) with 2 mM of DMAP for four minutes;
- 2- Move the oocytes into culture media with 2 mM of DMAP and 200 microM of Roscovitin for three hours;
- 3- Rinse four times and place in culture.

3. Activation by exposure to Ionomycin followed by cytochalasin and cycloheximide.

- 1- Place oocytes in Ionomycin (5 microM) for four minutes;
- 2- Move oocytes to culture media containing $5\ \mu\text{g/ml}$ of cytochalasin B and $5\ \mu\text{g/ml}$ of cycloheximide for five hours;
- 3- Rinse four times and place in culture.

4. Activation by electrical pulses

- 1- Place eggs in mannitol media containing $100\ \mu\text{M}$ CaCl_2 ;
- 2- Deliver three pulses of $1.0\ \text{kVcm}^{-1}$ for $20\ \mu\text{sec}$, each pulse 22 minutes apart;
- 3- Move oocytes to culture media containing $5\ \mu\text{g/ml}$ of cytochalasin B for three hours.

5. Activation by exposure with ethanol followed by cytochalasin and cycloheximide

- 1- Place oocytes in 7% ethanol for one minute;
- 2- Move oocytes to culture media containing $5\ \mu\text{g/ml}$ of cytochalasin B and $5\ \mu\text{g/ml}$ of cycloheximide for five hours;
- 3- Rinse four times and place in culture.

6. Activation by microinjection of adenophostine

- 1- Inject oocytes with 10 to 12 picoliters of a solution containing $10\ \mu\text{M}$ of adenophostine;
- 2- Put oocytes in culture.

7. Activation by microinjection of sperm factor

- 1- Inject oocytes with 10 to 12 picoliters of sperm factor isolated, e.g., from primates, pigs, bovine, sheep, goats, horses, mice, rats, rabbits or hamsters;
- 2- Put oocyte in culture.

8. **Activation by microinjection of recombinant sperm factor.**
9. **Activation by Exposure to DMAP followed by Cycloheximide and Cytochalasin B**

5 Place oocytes or NT units, typically about 22 to 28 hours post maturation in about 2 mM DMAP for about one hour, followed by incubation for about two to twelve hours, preferably about eight hours, in 5 μ g/ml of cytochalasin B and 20 μ g/ml cycloheximide.

10 The above activation protocols are exemplary of protocols used for nuclear transfer procedures, e.g., those including the use of primate or human donor cells or oocytes. However, the above activation protocols may be used when either or both the donor cell and nucleus is of ungulate origin, e.g., a sheep, buffalo, horse, goat, bovine, pig and/or wherein the oocyte is of ungulate origin, e.g., sheet, pig, buffalo, horse, goat, bovine, etc., as well as for other species.

15 As noted, activation may be effected before, simultaneous, or after nuclear transfer. In general, activation will be effected about 40 hours prior to nuclear transfer and fusion to about 40 hours after nuclear transfer and fusion, more preferably about 24 hours before to about 24 hours after nuclear transfer and fusion, and most preferably from about 4 to 9 hours before nuclear transfer and fusion to about 4 to 9 hours after nuclear
20 transfer and fusion. Activation is preferably effected after or proximate to *in vitro* or *in vivo* maturation of the oocyte, e.g., approximately simultaneous or within about 40 hours of maturation, more preferably within about 24 hours of maturation.

Activated NT units may be cultured in a suitable *in vitro* culture medium until the generation of embryonic or stem-like cells and cell colonies. Culture media suitable for culturing and maturation of embryos are well known in the art. Examples of known media, which may be used for bovine embryo culture and maintenance, include Ham's F-10 + 10% fetal calf serum (FCS), Tissue Culture Medium-199 (TCM-199) + 10% fetal calf serum, Tyrodes-Albumin-Lactate-Pyruvate (TALP), Dulbecco's Phosphate Buffered Saline (PBS), Eagle's and Whitten's media. One of the most common media used for the collection and maturation of oocytes is TCM-199, and 1 to 20% serum supplement including fetal calf serum, newborn serum, estrual cow serum, lamb serum or steer serum. A preferred maintenance medium includes TCM-199 with Earl salts, 10% fetal calf serum, 0.2 Ma pyruvate and 50 µg/ml gentamicin sulphate. Any of the above may also involve co-culture with a variety of cell types such as granulosa cells, oviduct cells, BRL cells and uterine cells and STO cells.

In particular, human epithelial cells of the endometrium secrete leukemia inhibitory factor (LIF) during the preimplantation and implantation period. Therefore, the addition of LIF to the culture medium could be of importance in enhancing the *in vitro* development of the reconstructed embryos. The use of LIF for embryonic or stem-like cell cultures has been described in U.S. Patent 5,712,156, which is herein incorporated by reference.

Another maintenance medium is described in U.S. Patent 5,096,822 to Rosenkrans, Jr. et al., which is incorporated herein by reference. This embryo medium, named CR1,

contains the nutritional substances necessary to support an embryo. CR1 contains hemicalcium L-lactate in amounts ranging from 1.0 mM to 10 mM, preferably 1.0 mM to 5.0 mM. Hemicalcium L-lactate is L-lactate with a hemicalcium salt incorporated therein.

5 Also, suitable culture medium for maintaining human embryonic cells in culture are discussed in Thomson et al., *Science*, 282:1145-1147 (1998) and *Proc. Natl. Acad. Sci., USA*, 92:7844-7848 (1995). Further, the NT units may be cultured in biological fluids such as peritoneal fluid, amniotic fluid, vitreous/aqueous humor and lymph fluid.

Afterward, the cultured NT unit or units are preferably washed and then placed in a suitable media, e.g., the media identified above, such as CR1aa medium, Ham's F-10, Tissue Culture Media -199 (TCM-199). Tyrodes-Albumin-Lactate-Pyruvate (TALP) Dulbecco's Phosphate Buffered Saline (PBS), Eagle's or Whitten's, preferably containing about 10% FCS. Such culturing will preferably be effected in well plates which contain a suitable confluent feeder layer. Suitable feeder layers include, by way of example, 10 fibroblasts and epithelial cells, e.g., fibroblasts and uterine epithelial cells derived from ungulates, chicken fibroblasts, murine (e.g., mouse or rat) fibroblasts, STO and SI-m220 feeder cell lines, and BRL cells.

In the preferred embodiment, the feeder cells will comprise mouse embryonic fibroblasts. Means for preparation of a suitable fibroblast feeder layer are described in the example which follows and is well within the skill of the ordinary artisan.

The NT units are cultured on the feeder layer until the NT units reach a size suitable for obtaining cells which may be used to produce embryonic stem-like cells or cell colonies. Preferably, these NT units will be cultured until they reach a size of at least about 2 to 400 cells, more preferably about 4 to 128 cells, and most preferably at least about 50 cells. The culturing will be effected under suitable conditions, i.e., about 38.5°C and 5% CO₂, with the culture medium changed in order to optimize growth typically about every 2-5 days, preferably about every 3 days.

In the case of human cell/enucleated bovine oocyte derived NT units, sufficient cells to produce an ES cell colony, typically on the order of about 50 cells, will be obtained about 12 days after initiation of oocyte activation. However, this may vary dependent upon the particular cell used as the nuclear donor, the species of the particular oocyte, and culturing conditions. One skilled in the art can readily ascertain visually when a desired sufficient number of cells has been obtained based on the morphology of the cultured NT units.

In the case of human/human nuclear transfer embryos, or other embryos produced using non-human primate donor or oocyte, it may be advantageous to use culture medium known to be useful for maintaining human and other primate cells in tissue culture. Examples of a culture media suitable for human embryo culture include the medium reported in Jones et al, *Human Reprod.*, 13(1):169-177 (1998), the P1-catalog #99242 medium, and the P-1 catalog #99292 medium, both available from Irvine Scientific, Santa

Ana, California, and those used by Thomson et al. (1998) and (1995), which references are incorporated by reference in their entirety.

Another preferred medium comprises ACM + uridine + glucose + 1000 IU of LIF.

Still other examples of suitable media include naturally occurring biological fluids
5 such as peritoneal fluid, vitreous /aqueous humor, amniotic fluid, and lymph fluid.

As discussed above, the cells used in the present invention will preferably
comprise mammalian somatic cells, most preferably cells derived from an actively
proliferating (non-quiescent) mammalian cell culture. In an especially preferred
embodiment, the donor cell will be genetically modified by the addition, deletion or
substitution of a desired DNA sequence. For example, the donor cell, e.g., a keratinocyte
10 or fibroblast, e.g., of human, primate or bovine origin, may be transfected or transformed
with a DNA construct that provides for the expression of a desired gene product, e.g.,
therapeutic polypeptide. Examples thereof include lymphokines, e.g., IGF-I, IGF-II,
interferons, colony stimulating factors, connective tissue polypeptides such as collagens,
15 genetic factors, clotting factors, enzymes, enzyme inhibitors, etc. Also, desired genes
may be "knocked in" or "knocked out" by homologous recombination.

Also, as discussed above, the donor cells may be modified prior to nuclear transfer,
e.g., to effect impaired cell lineage development, enhanced embryonic development
and/or inhibition of apoptosis. Examples of desirable modifications are discussed further
20 below.

One aspect of the invention will involve genetic modification of the donor cell, e.g., a human cell, such that it is lineage deficient and therefore when it is used for nuclear transfer it will be unable to give rise to a viable offspring. This is desirable especially in the context of human nuclear transfer embryos, wherein for ethical reasons, production of a viable embryo may be an unwanted outcome. This can be effected by genetically engineering a human cell such that it is incapable of differentiating into specific cell lineages when used for nuclear transfer. In particular, cells may be genetically modified such that when used as nuclear transfer donors the resultant "embryos" do not contain or substantially lack at least one of mesoderm, endoderm or ectoderm tissue.

This can be accomplished by, e.g., knocking out or impairing the expression of one or more mesoderm, endoderm or ectoderm specific genes. Examples thereof include:

Mesoderm: SRF, MESP-1, HNF-4, beta-I integrin, MSD;

Endoderm: GATA-6, GATA-4;

Ectoderm: RNA helicase A, H beta 58.

The above list is intended to be exemplary and non-exhaustive of known genes which are involved in the development of mesoderm, endoderm and ectoderm. The generation of mesoderm deficient, endoderm deficient and ectoderm deficient cells and embryos has been previously reported in the literature. See, e.g., Arsenian et al, *EMBO J.*, Vol. 17(2):6289-6299 (1998); Saga Y, *Mech. Dev.*, Vol. 75(1-2):53-66 (1998); Holdener et al, *Development*, Vol. 120(5):1355-1346 (1994); Chen et al, *Genes Dev.* Vol. 8(20):2466-2477 (1994); Rohwedel et al, *Dev. Biol.*, 201(2):167-189 (1998) (mesoderm);

Morrissey et al, *Genes, Dev.*, Vol. 12(22):3579-3590 (1998); Soudais et al, *Development*, Vol. 121(11):3877-3888 (1995) (endoderm); and Lee et al, *Proc. Natl. Acad. Sci. USA*, Vol. 95:(23):13709-13713 (1998); and Radice et al, *Development*, Vol. 111(3):801-811 (1991) (ectoderm).

5 In general, a desired somatic cell, e.g., a human keratinocyte, epithelial cell or fibroblast, will be genetically engineered such that one or more genes specific to particular cell lineages are "knocked out" and/or the expression of such genes significantly impaired. This may be effected by known methods, e.g., homologous recombination. A preferred genetic system for effecting "knock-out" of desired genes is disclosed by Capecchi et al, U.S. Patents 5,631,153 and 5,464,764, which reports
10 positive-negative selection (PNS) vectors that enable targeted modification of DNA sequences in a desired mammalian genome. Such genetic modification will result in a cell that is incapable of differentiating into a particular cell lineage when used as a nuclear transfer donor.

15 This genetically modified cell will be used to produce a lineage-defective nuclear transfer embryo, i.e., that does not develop at least one of a functional mesoderm, endoderm or ectoderm. Thereby, the resultant embryos, even if implanted, e.g., into a human uterus, would not give rise to a viable offspring. However, the ES cells that result from such nuclear transfer will still be useful in that they will produce cells of the one or
20 two remaining non-impaired lineage. For example, an ectoderm deficient human nuclear transfer embryo will still give rise to mesoderm and endoderm derived differentiated

cells. An ectoderm deficient cell can be produced by deletion and/or impairment of one or both of RNA helicase A or H beta 58 genes.

These lineage deficient donor cells may also be genetically modified to express another desired DNA sequence. Thus, the genetically modified donor cell will give rise to a lineage-deficient blastocyst which, when plated, will differentiate into at most two of the embryonic germ layers.

Alternatively, the donor cell can be modified such that it is "immortal". This can be achieved by expressing anti-sense or ribozyme telomerase genes. This can be effected by known genetic methods that will provide for expression of antisense DNA or ribozymes, or by gene knockout. These "immortal" cells, when used for nuclear transfer, will not be capable of differentiating into viable offspring.

Another preferred embodiment of the present invention is the production of nuclear transfer embryos that grow more efficiently in tissue culture. This is advantageous in that it should reduce the requisite time and necessary fusions to produce ES cells and/or offspring (if the blastocysts are to be implanted into a female surrogate). This is desirable also because it has been observed that blastocysts and ES cells resulting from nuclear transfer may have impaired development potential. While these problems may often be alleviated by alteration of tissue culture conditions, an alternative solution is to enhance embryonic development by enhancing expression of genes involved in embryonic development.

For example, it has been reported that the gene products of the Ped type, which are members of the MHC I family, are of significant importance to embryonic development. More specifically, it has been reported in the case of mouse preimplantation embryos that the Q7 and Q9 genes are responsible for the "fast growth" phenotype. Therefore, it is anticipated that introduction of DNAs that provide for the expression of these and related genes, or their human or other mammalian counterparts into donor cells, will give rise to nuclear transfer embryos that grow more quickly. This is particularly desirable in the context of cross-species nuclear transfer embryos which may develop less efficiently in tissue culture than nuclear transfer embryos produced by fusion of cells or nuclei of the same species.

Preferably, a DNA construct containing the Q7 and/or Q9 gene will be introduced into donor somatic cells prior to nuclear transfer. For example, an expression construct can be constructed containing a strong constitutive mammalian promoter operably linked to the Q7 and/or Q9 genes, an IRES, one or more suitable selectable markers, e.g., neomycin, ADA, DHFR, and a poly-A sequence, e.g., bGH polyA sequence. Also, it may be advantageous to further enhance Q7 and Q9 gene expression by the inclusion of insulates. It is anticipated that these genes will be expressed early on in blastocyst development as these genes are highly conserved in different species, e.g., bovines, goats, porcine, dogs, cats, and humans. Also, it is anticipated that donor cells can be engineered to affect other genes that enhance embryonic development. Thus, these genetically

modified donor cells should produce blastocysts and preimplantation stage embryos more efficiently.

Still another aspect of the invention involves the construction of donor cells that are resistant to apoptosis, i.e., programmed cell death. It has been reported in the literature that cell death related genes are present in preimplantation stage embryos. (Adams et al, *Science*, 281(5381):1322-1326 (1998)). Genes reported to induce apoptosis include, e.g., Bad, Bok, BH3, Bik, Hrk, BNIP3, Bim_L, Bad, Bid, and EGL-1. By contrast, genes that reportedly protect cells from programmed cell death include, by way of example, Bcl-XL, Bcl-w, Mcl-1, A1, Nr-13, BHRF-1, LMW5-HL, ORF16, Ks-Bcl-2, E1B-19K, and CED-9.

Thus, donor cells can be constructed wherein genes that induce apoptosis are "knocked out" or wherein the expression of genes that protect the cells from apoptosis is enhanced or turned on during embryonic development.

For example, this can be effected by introducing a DNA construct that provides for regulated expression of such protective genes, e.g., Bcl-2 or related genes during embryonic development. Thereby, the gene can be "turned on" by culturing the embryo under specific growth conditions. Alternatively, it can be linked to a constitutive promoter.

More specifically, a DNA construct containing a Bcl-2 gene operably linked to a regulatable or constitutive promoter, e.g., PGK, SV40, CMV, ubiquitin, or beta-actin, an

IRES, a suitable selectable marker, and a poly-A sequence can be constructed and introduced into a desired donor mammalian cell, e.g., human keratinocyte or fibroblast.

These donor cells, when used to produce nuclear transfer embryos, should be resistant to apoptosis and thereby differentiate more efficiently in tissue culture. Thereby, the speed and/or number of suitable preimplantation embryos produced by nuclear transfer can be increased.

Another means of accomplishing the same result is to impair the expression of one or more genes that induce apoptosis. This will be effected by knock-out or by the use of antisense or ribozymes against genes that are expressed in and which induce apoptosis early on in embryonic development. Examples thereof are identified above. Cell death genes that may be expressed in the antisense orientation include BAX, Apaf-1, and capsases. Additionally, a transgene may be introduced that encodes for methylase or demethylase in the sense or antisense orientation. DNAs that encode methylase and demethylase enzymes are well known in the art. Still alternatively, donor cells may be constructed containing both modifications, i.e., impairment of apoptosis-inducing genes and enhanced expression of genes that impede or prevent apoptosis. The construction and selection of genes that affect apoptosis, and cell lines that express such genes, is disclosed in U.S. Patent No. 5,646,008, which patent is incorporated by reference herein. Many DNAs that promote or inhibit apoptosis have been reported and are the subject of numerous patents.

Another means of enhancing cloning efficiency is to select cells of a particular cell cycle stage as the donor cell. It has been reported that this can have significant effects on nuclear transfer efficiency. (Barnes et al, *Mol. Reprod. Devel.*, 36(1):33-41 (1993). Different methods for selecting cells of a particular cell cycle stage have been reported and include serum starvation (Campbell et al, *Nature*, 380:64-66 (1996); Wilmut et al, *Nature*, 385:810-813 (1997), and chemical synchronization (Urbani et al, *Exp. Cell Res.*, 219(1):159-168 (1995). For example, a particular cyclin DNA may be operably linked to a regulatory sequence, together with a detectable marker, e.g., green fluorescent protein (GFP), followed by the cyclin destruction box, and optionally insulation sequences to enhance cyclin and marker protein expression. Thereby, cells of a desired cell cycle can be easily visually detected and selected for use as a nuclear transfer donor. An example thereof is the cyclin D1 gene in order to select for cells that are in G1. However, any cyclin gene should be suitable for use in the claimed invention. (See, e.g., King et al, *Mol. Biol. Cell*, Vol. 7(9):1343-1357 (1996)).

However, a less invasive or more efficient method for producing cells of a desired cell cycle stage are needed. It is anticipated that this can be effected by genetically modifying donor cells such that they express specific cyclins under detectable conditions. Thereby, cells of a specific cell cycle can be readily discerned from other cell cycles.

Cyclins are proteins that are expressed only during specific stages of the cell cycle. They include cyclin D1, D2 and D3 in G1 phase, cyclin B1 and B2 in G2/M phase and cyclin E, A and H in S phase. These proteins are easily translated and destroyed in the

cytogolcytosol. This "transient" expression of such proteins is attributable in part to the presence of a "destruction box", which is a short amino acid sequence that is part of the protein that functions as a tag to direct the prompt destruction of these proteins via the ubiquitin pathway. (Adams et al, *Science*, 281 (5321):1322-1326 (1998)).

5 In the present invention, donor cells will be constructed that express one or more of such cyclin genes under easily detectable conditions, preferably visualizable, e.g., by the use of a fluorescent label. For example, a particular cyclin DNA may be operably linked to a regulatory sequence, together with a detectable marker, e.g., green fluorescent protein (GFP), followed by the cyclin destruction box, and optionally insulation sequences to enhance cyclin and/or marker protein expression. Thereby, cells of a
10 desired cell cycle can be easily visually detected and selected for use as a nuclear transfer donor. An example thereof is the cyclin D1 gene which can be used to select for cells that are in G1. However, any cyclin gene should be suitable for use in the claimed invention. (See, e.g., King et al, *Mol. Biol. Cell*, Vol. 7(9):1343-1357 (1996)).

15 As discussed, the present invention provides different methods for enhancing nuclear transfer efficiency, preferably a cross-species nuclear transfer process. While the present inventors have demonstrated that nuclei or cells of one species when inserted or fused with an enucleated oocyte of a different species can give rise to nuclear transfer embryos that produce blastocysts, which embryos can give rise to ES cell lines, the
20 efficiency of such process is quite low. Therefore, many fusions typically need to be effected to produce a blastocyst the cells of which may be cultured to produce ES cells

and ES cell lines. Yet another means for enhancing the development of nuclear transfer embryos *in vitro* is by optimizing culture conditions. One means of achieving this result will be to culture NT embryos under conditions impede apoptosis. With respect to this embodiment of the invention, it has been found that proteases such as capsases can cause oocyte death by apoptosis similar to other cell types. (See, Jurisicosva et al, *Mol. Reprod. Devel.*, 51(3):243-253 (1998).)

It is anticipated that blastocyst development will be enhanced by including in culture media used for nuclear transfer and to maintain blastocysts or culture preimplantation stage embryos one or more capsase inhibitors. Such inhibitors include by way of example capsase-4 inhibitor I, capsase-3 inhibitor I, capsase-6 inhibitor II, capsase-9 inhibitor II, and capsase-1 inhibitor I. The amount thereof will be an amount effective to inhibit apoptosis, e.g., 0.00001 to 5.0% by weight of medium; more preferably 0.01% to 1.0% by weight of medium. Thus, the foregoing methods may be used to increase the efficiency of nuclear transfer by enhancing subsequent blastocyst and embryo development in tissue culture.

After NT units of the desired size are obtained, the cells are mechanically removed from the zone and are then used to produce embryonic or stem-like cells and cell lines. This is preferably effected by taking the clump of cells which comprise the NT unit, which typically will contain at least about 50 cells, washing such cells, and plating the cells onto a feeder layer, e.g., irradiated fibroblast cells. Typically, the cells used to obtain the stem-like cells or cell colonies will be obtained from the inner most portion of

the cultured NT unit which is preferably at least 50 cells in size. However, NT units of smaller or greater cell numbers as well as cells from other portions of the NT unit may also be used to obtain ES-like cells and cell colonies.

It is further envisioned that a longer exposure of donor cell DNA to the oocyte's cytosol may facilitate the dedifferentiation process. This can be accomplished by re-cloning, i.e., by taking blastomeres from a reconstructed embryo and fusing them with a new enucleated oocyte. Alternatively, the donor cell may be fused with an enucleated oocyte and four to six hours later, without activation, chromosomes removed and fused with a younger oocyte. Activation would occur thereafter.

The cells are maintained in the feeder layer in a suitable growth medium, e.g., alpha MEM supplemented with 10% FCS and 0.1 mM beta-mercaptoethanol (Sigma) and L-glutamine. The growth medium is changed as often as necessary to optimize growth, e.g., about every 2-3 days.

This culturing process results in the formation of embryonic or stem-like cells or cell lines. In the case of human cell/bovine oocyte derived NT embryos, colonies are observed by about the second day of culturing in the alpha MEM medium. However, this time may vary dependent upon the particular nuclear donor cell, specific oocyte and culturing conditions. One skilled in the art can vary the culturing conditions as desired to optimize growth of the particular embryonic or stem-like cells. Other suitable media are disclosed herein.

5 The embryonic or stem-like cells and cell colonies obtained will typically exhibit an appearance similar to embryonic or stem-like cells of the species used as the nuclear cell donor rather than the species of the donor oocyte. For example, in the case of embryonic or stem-like cells obtained by the transfer of a human nuclear donor cell into an enucleated bovine oocyte, the cells exhibit a morphology more similar to mouse embryonic stem cells than bovine ES-like cells.

10 More specifically, the individual cells of the human ES-line cell colony are not well defined, and the perimeter of the colony is refractive and smooth in appearance. Further, the cell colony has a longer cell doubling time, about twice that of mouse ES cells. Also, unlike bovine and porcine derived ES cells, the colony does not possess an epithelial-like appearance.

15 As discussed above, it has been reported by Thomson, in U.S. Patent 5,843,780, that primate stem cells are SSEA-1 (-), SSEA-4 (+), TRA-1-60 (+), TRA-1-81 (+) and alkaline phosphatase (+). It is anticipated that human and primate ES cells produced according to the present methods will exhibit similar or identical marker expression.

20 Alternatively, that such cells are actual human or primate embryonic stem cells will be confirmed based on their capability of giving rise to all of mesoderm, ectoderm and endoderm tissues. This will be demonstrated by culturing ES cells produced according to the invention under appropriate conditions, e.g., as disclosed by Thomsen, U.S. Patent 5,843,780, incorporated by reference in its entirety herein. Alternatively, the fact that the cells produced according to the invention are pluripotent will be confirmed

by injecting such cells into an animal, e.g., a SCID mouse, or large agricultural animal, and thereafter obtaining tissues that result from said implanted cells. These implanted ES cells should give rise to all different types of differentiated tissues, i.e., mesoderm, ectoderm, and endodermal tissues.

5 The resultant embryonic or stem-like cells and cell lines, preferably human embryonic or stem-like cells and cell lines, have numerous therapeutic and diagnostic applications. Most especially, such embryonic or stem-like cells may be used for cell transplantation therapies. Human embryonic or stem-like cells have application in the treatment of numerous disease conditions.

10 In this regard, it is known that mouse embryonic stem (ES) cells are capable of differentiating into almost any cell type, e.g., hematopoietic stem cells. Therefore, human embryonic or stem-like cells produced according to the invention should possess similar differentiation capacity. The embryonic or stem-like cells according to the invention will be induced to differentiate to obtain the desired cell types according to known methods.

15 For example, the subject human embryonic or stem-like cells may be induced to differentiate into hematopoietic stem cells, muscle cells, cardiac muscle cells, liver cells, cartilage cells, epithelial cells, urinary tract cells, etc., by culturing such cells in differentiation medium and under conditions which provide for cell differentiation. Medium and methods which result in the differentiation of embryonic stem cells are

20 known in the art as are suitable culturing conditions.

For example, Palacios et al, *Proc. Natl. Acad. Sci., USA*, 92:7530-7537 (1995) teaches the production of hematopoietic stem cells from an embryonic cell line by subjecting stem cells to an induction procedure comprising initially culturing aggregates of such cells in a suspension culture medium lacking retinoic acid followed by culturing in the same medium containing retinoic acid, followed by transferral of cell aggregates to a substrate which provides for cell attachment.

Moreover, Pedersen, *J. Reprod. Fertil. Dev.*, 6:543-552 (1994) is a review article which references numerous articles disclosing methods for *in vitro* differentiation of embryonic stem cells to produce various differentiated cell types including hematopoietic cells, muscle, cardiac muscle, nerve cells, among others.

Further, Bain et al, *Dev. Biol.*, 168:342-357 (1995) teaches *in vitro* differentiation of embryonic stem cells to produce neural cells which possess neuronal properties. These references are exemplary of reported methods for obtaining differentiated cells from embryonic or stem-like cells. These references and in particular the disclosures therein relating to methods for differentiating embryonic stem cells are incorporated by reference in their entirety herein.

Thus, using known methods and culture medium, one skilled in the art may culture the subject embryonic or stem-like cells to obtain desired differentiated cell types, e.g., neural cells, muscle cells, hematopoietic cells, etc. In addition, the use of inducible Bcl-2 or Bcl-xl might be useful for enhancing *in vitro* development of specific cell lineages. *In vivo*, Bcl-2 prevents many, but not all, forms of apoptotic cell death that occur during

lymphoid and neural development. A thorough discussion of how Bcl-2 expression might be used to inhibit apoptosis of relevant cell lineages following transfection of donor cells is disclosed in U.S. Patent No. 5,646,008, which is herein incorporated by reference.

The subject embryonic or stem-like cells may be used to obtain any desired differentiated cell type. Therapeutic usages of such differentiated human cells are unparalleled. For example, human hematopoietic stem cells may be used in medical treatments requiring bone marrow transplantation. Such procedures are used to treat many diseases, e.g., late stage cancers such as ovarian cancer and leukemia, as well as diseases that compromise the immune system, such as AIDS. Hematopoietic stem cells can be obtained, e.g., by fusing adult somatic cells of a cancer or AIDS patient, e.g., epithelial cells or lymphocytes with an enucleated oocyte, e.g., bovine oocyte, obtaining embryonic or stem-like cells as described above, and culturing such cells under conditions which favor differentiation, until hematopoietic stem cells are obtained. Such hematopoietic cells may be used in the treatment of diseases including cancer and AIDS.

Alternatively, adult somatic cells from a patient with a neurological disorder may be fused with an enucleated animal oocyte, e.g., a primate or bovine oocyte, human embryonic or stem-like cells obtained therefrom, and such cells cultured under differentiation conditions to produce neural cell lines. Specific diseases treatable by transplantation of such human neural cells include, by way of example, Parkinson's disease, Alzheimer's disease, ALS and cerebral palsy, among others. In the specific case of Parkinson's disease, it has been demonstrated that transplanted fetal brain neural cells

make the proper connections with surrounding cells and produce dopamine. This can result in long-term reversal of Parkinson's disease symptoms.

To allow for specific selection of differentiated cells, donor cells may be transfected with selectable markers expressed via inducible promoters, thereby permitting selection or enrichment of particular cell lineages when differentiation is induced. For example, CD34-neo may be used for selection of hematopoietic cells, Pw1-neo for muscle cells, Mash-1-neo for sympathetic neurons, Mal-neo for human CNS neurons of the grey matter of the cerebral cortex, etc. Also, the use of matrices to facilitate tissue development may be beneficial.

The great advantage of the subject invention is that it provides an essentially limitless supply of isogenic or syngenic human cells suitable for transplantation. Therefore, it will obviate the significant problem associated with current transplantation methods, i.e., rejection of the transplanted tissue which may occur because of host-vs-graft or graft-vs-host rejection. Conventionally, rejection is prevented or reduced by the administration of anti-rejection drugs such as cyclosporin. However, such drugs have significant adverse side-effects, e.g., immunosuppression, carcinogenic properties, as well as being very expensive. The present invention should eliminate, or at least greatly reduce, the need for anti-rejection drugs, such as cyclosporine, imulan, FK-506, glucocorticoids, rapamycin, and derivatives thereof.

Other diseases and conditions treatable by isogenic cell therapy include, by way of example, spinal cord injuries, multiple sclerosis, muscular dystrophy, diabetes, liver diseases, i.e., hypercholesterolemia, heart diseases, cartilage replacement, burns, foot ulcers, gastrointestinal diseases, vascular diseases, kidney disease, urinary tract disease, and aging related diseases and conditions.

Also, human embryonic or stem-like cells produced according to the invention may be used to produce genetically engineered or transgenic human differentiated cells. Essentially, this will be effected by introducing a desired gene or genes, which may be heterologous, or removing all or part of an endogenous gene or genes of human embryonic or stem-like cells produced according to the invention, and allowing such cells to differentiate into the desired cell type. A preferred method for achieving such modification is by homologous recombination because such technique can be used to insert, delete or modify a gene or genes at a specific site or sites in the stem-like cell genome.

This methodology can be used to replace defective genes, e.g., defective immune system genes, cystic fibrosis genes, or to introduce genes which result in the expression of therapeutically beneficial proteins such as growth factors, lymphokines, cytokines, enzymes, etc. For example, the gene encoding brain derived growth factor may be introduced into human embryonic or stem-like cells, the cells differentiated into neural cells and the cells transplanted into a Parkinson's patient to retard the loss of neural cells during such disease.

Previously, cell types transfected with BDNF varied from primary cells to immortalized cell lines, either neural or non-neural (myoblast and fibroblast) derived cells. For example, astrocytes have been transfected with BDNF gene using retroviral vectors, and the cells grafted into a rat model of Parkinson's disease (Yoshimoto et al., *Brain Research*, 691:25-36, (1995)).

This *ex vivo* therapy reduced Parkinson's-like symptoms in the rats up to 45% 32 days after transfer. Also, the tyrosine hydroxylase gene has been placed into astrocytes with similar results (Lundberg et al., *Develop. Neurol.*, 139:39-53 (1996) and references cited therein).

However, such *ex vivo* systems have problems. In particular, retroviral vectors currently used are down-regulated *in vivo* and the transgene is only transiently expressed (review by Mulligan, *Science*, 260:926-932 (1993)). Also, such studies used primary cells, astrocytes, which have finite life span and replicate slowly. Such properties adversely affect the rate of transfection and impede selection of stably transfected cells. Moreover, it is almost impossible to propagate a large population of gene targeted primary cells to be used in homologous recombination techniques.

By contrast, the difficulties associated with retroviral systems should be eliminated by the use of human embryonic or stem-like cells. It has been demonstrated previously by the subject assignee that cattle and pig embryonic cell lines can be transfected and selected for stable integration of heterologous DNA. Such methods are described in commonly assigned U.S. Patent No. 5,905,042, incorporated by reference in its entirety.

Therefore, using such methods or other known methods, desired genes may be introduced into the subject human embryonic or stem-like cells, and the cells differentiated into desired cell types, e.g., hematopoietic cells, neural cells, pancreatic cells, cartilage cells, etc.

5 Genes which may be introduced into the subject embryonic or stem-like cells include, by way of example, epidermal growth factor, basic fibroblast growth factor, glial derived neurotrophic growth factor, insulin-like growth factor (I and II), neurotrophin-3, neurotrophin-4/5, ciliary neurotrophic factor, AFT-1, cytokine genes (interleukins, interferons, colony stimulating factors, tumor necrosis factors (alpha and beta), etc.),
10 genes encoding therapeutic enzymes, collagen, human serum albumin, etc.

In addition, it is also possible to use one of the negative selection systems now known in the art for eliminating therapeutic cells from a patient if necessary. For example, donor cells transfected with the thymidine kinase (TK) gene will lead to the production of embryonic cells containing the TK gene. Differentiation of these cells will
15 lead to the isolation of therapeutic cells of interest which also express the TK gene. Such cells may be selectively eliminated at any time from a patient upon gancyclovir administration. Such a negative selection system is described in U.S. Patent No. 5,698,446, and is herein incorporated by reference.

20 The subject embryonic or stem-like cells, preferably human cells, also may be used as an *in vitro* model of differentiation, in particular for the study of genes which are involved in the regulation of early development.

Also, differentiated cell tissues and organs using the subject embryonic or stem-like cells may be used in drug studies.

Further, the subject cells may be used to express recombinant DNAs.

Still further, the subject embryonic or stem-like cells may be used as nuclear donors for the production of other embryonic or stem-like cells and cell colonies.

Also, cultured inner cell mass, or stem cells, produced according to the invention may be introduced into animals, e.g., SCID mice, cows, pigs, e.g., under the renal capsule or intramuscularly and used to produce a teratoma therein. This teratoma can be used to derive different tissue types. Also, the inner cell mass produced by X-species nuclear transfer may be introduced together with a biodegradable, biocompatible polymer matrix that provides for the formation of 3-dimensional tissues. After tissue formation, the polymer degrades, ideally just leaving the donor tissue, e.g., cardiac, pancreatic, neural, lung, liver. In some instances, it may be advantageous to include growth factors and proteins that promote angiogenesis. Alternatively, the formation of tissues can be effected totally *in vitro*, with appropriate culture media and conditions, growth factors, and biodegradable polymer matrices.

In order to more clearly describe the subject invention, the following examples are provided.

EXAMPLE 1

MATERIALS AND METHODS

Donor Cells for Nuclear Transfer

Epithelial cells were lightly scraped from the inside of the mouth of a consenting adult with a standard glass slide. The cells were washed off the slide into a petri dish containing phosphate buffered saline without Ca or Mg. The cells were pipetted through a small-bore pipette to break up cell clumps into a single cell suspension. The cells were then transferred into a microdrop of TL-HEPES medium containing 10% fetal calf serum (FCS) under oil for nuclear transfer into enucleated cattle oocytes.

Nuclear Transfer Procedures

Basic nuclear transfer procedures have been described previously. Briefly, after slaughterhouse oocytes were matured *in vitro* the oocytes were stripped of cumulus cells and enucleated with a beveled micropipette at approximately 18 hours post maturation (hpm). Enucleation was confirmed in TL-HEPES medium plus bisbenzimidazole (Hoechst 33342, 3 µg/ml; Sigma). Individual donor cells were then placed into the perivitelline space of the recipient oocyte. The bovine oocyte cytoplasm and the donor nucleus (NT unit) are fused together using electrofusion techniques. One fusion pulse consisting of 90 V for 15 µsec was applied to the NT unit. This occurred at 24 hours post-initiation of maturation (hpm) of the oocytes. The NT units were placed in CR1aa medium until 28 hpm.

The procedure used to artificially activate oocytes has been described elsewhere. NT unit activation was at 28 hpm. A brief description of the activation procedure is as follows: NT units were exposed for four min to ionomycin (5 µM; CalBiochem, La Jolla, CA) in TL-HEPES supplemented with 1 mg/ml BSA and then washed for five min in TL-

HEPES supplemented with 30 mg/ml BSA. The NT units were then transferred into a microdrop of CR1aa culture medium containing 0.2 mM DMAP (Sigma) and cultured at 38.5°C 5% CO₂ for four to five hours. The NT units were washed and then placed in a CR1aa medium plus 10% FCS and 6 mg/ml BSA in four well plates containing a confluent feeder layer of mouse embryonic fibroblasts (described below). The NT units were cultured for three more days at 38.5°C and 5% CO₂. The culture medium was changed every three days until day 12 after the time of activation. At this time NT units reaching the desired cell number, i.e., about 50 cell number, were mechanically removed from the zona and used to produce embryonic cell lines. A photograph of an NT unit obtained as described above is contained in Figure 1.

Fibroblast feeder layer

Primary cultures of embryonic fibroblasts were obtained from 14-16 day old murine fetuses. After the head, liver, heart and alimentary tract were aseptically removed, the embryos were minced and incubated for 30 minutes at 37°C in pre-warmed trypsin EDTA solution (0.05% trypsin/0.02% EDTA; GIBCO, Grand Island, NY). Fibroblast cells were plated in tissue culture flasks and cultured in alpha-MEM medium (BioWhittaker, Walkersville, MD) supplemented with 10% fetal calf serum (FCS) (Hyclone, Logan, UT), penicillin (100 IU/ml) and streptomycin (50 µl/ml). Three to four days after passage, embryonic fibroblasts, in 35 x 10 Nunc culture dishes (Baxter Scientific, McGaw Park, IL), were irradiated. The irradiated fibroblasts were grown and

maintained in a humidified atmosphere with 5% CO₂ in air at 37°C. The culture plates which had a uniform monolayer of cells were then used to culture embryonic cell lines.

Production of embryonic cell line.

NT unit cells obtained as described above were washed and plated directly onto irradiated feeder fibroblast cells. These cells included those of the inner portion of the NT unit. The cells were maintained in a growth medium consisting of alpha MEM supplemented with 10% FCS and 0.1 mM beta-mercaptoethanol (Sigma). Growth medium was exchanged every two to three days. The initial colony was observed by the second day of culture. The colony was propagated and exhibits a similar morphology to previously disclosed mouse embryonic stem (ES) cells. Individual cells within the colony are not well defined and the perimeter of the colony is refractile and smooth in appearance. The cell colony appears to have a slower cell doubling time than mouse ES cells. Also, unlike bovine and porcine derived ES cells, the colony does not have an epithelial appearance thus far. Figures 2 through 5 are photographs of ES-like cell colonies obtained as described, *supra*.

Production of Differentiated Human Cells

The human embryonic cells obtained are transferred to a differentiation medium and cultured until differentiated human cell types are obtained.

RESULTS

Table 1. Human cells as donor nuclei in NT unit production and development.

TABLE 1

Cell type	No. NT units made	No. NT units 2 cell stage (%)	No. NT units to 4 - 16 cell stage (%)	No. NT units to 16 - 400 cell stage (%)
lymphocytes	18	12 (67%)	3 (17%)	0
oral cavity epithelium	34	18 (53%)	3 (9%)	1 (3%)
adult fibroblasts	46	4 (9%)	12 (4 cell; 26%) 8 (8-16 cells; 17.4%)	---

The one NT unit that developed a structure having greater than 16 cells was plated down onto a fibroblast feeder layer. This structure was attached to the feeder layer and started to propagate forming a colony with a ES cell-like morphology (*See, e.g., Figure 2*). Moreover, although the 4 to 16 cell stage structures were not used to try and produce an ES cell colony, it has been previously shown that this stage is capable of producing ES or ES-like cell lines (mouse, Eistetter et al., *Devel. Growth and Differ.*, 31:275-282 (1989); *Bovine*, Stice et al., 1996)). Therefore, it is expected that 4 - 16 cell stage NT units should also give rise to embryonic or stem-like cells and cell colonies.

Also, similar results were obtained upon fusion of an adult human keratinocyte cell line with an enucleated bovine oocyte, which was cultured in media comprising ACM, uridine, glucose, and 1000 IU of LIF. Out of 50 reconstructed embryos, 22 cleaved and

one developed into a blastocyst at about day 12. This blastocyst was plated and the production of an ES cell line is ongoing.

EXAMPLE 2

A. Mitochondria Isolation Protocol from a Cell

This Example relates to isolation of mitochondria and use thereof to enhance the efficiency of cross-species nuclear transfer. The number of mitochondria per cell varies from cell line to cell line. For example, mouse L cells contain ~100 mitochondria per cell, whereas there are at least twice that number in HeLa cells. The cells are swollen in a hypotonic buffer and ruptured with a few strokes in a Dounce homogenizer using a tight-fitting pestle, and the mitochondria are isolated by differential centrifugation.

The solutions, tubes, and homogenizer should be pre-chilled on ice. All centrifugation steps are at 40°C. This protocol is based on starting with a washed cell pellet of 1-2 ml. The cell pellet is resuspended in 11 ml of ice-cold RSB and transferred to a 16 ml Dounce homogenizer.

RSB Buffer

RSB (A hypotonic buffer for swelling the tissue culture cells)

10 mM NaCl

1.5 mM MgCl₂

10 mM Tris-HCl, pH 7.5

MgCl₂

The cells are allowed to swell for five to ten minutes. The progress of the swelling is maintained using a phase contrast microscope. The swollen cells are replaced, preferably by several strokes with a pestle. Immediately after, 8 ml of 2.5x MS buffer are added to give a final concentration of 1x MS. The top of the homogenizer is then covered with Parafilm and mixed by inverting a couple of times.

2.5x MS Buffer

525 mM mannitol

175 Mm sucrose

12.5 mM ris-HCl, pH 7.5

215 mM EDTA pH 7.5

1x MS Buffer

210 mM mannitol

70 mM Sucrose

5 mM Tris-HCl, pH 7.5

1 mM EDTA, pH 7.5

MS Buffer is an iso-osmotic buffer to maintain the tonicity of the organelles and prevent agglutination.

Thereafter, the homogenate is transferred to a centrifuge tube for differential centrifugation. The homogenizer is rinsed with a small amount of MS buffer and added to the homogenate. The volume is brought to 30 ml with MS buffer. The homogenate is then centrifuged at 1300 g for five minutes to remove nuclei, unbroken cells, and large

membrane fragments. The supernatant is then poured into a clean centrifuge tube. The nuclear spin-down is repeated twice. The supernatant is then transferred to a clean centrifuge tube and a pellet containing the mitochondria is centrifuged at 17,000 g for 15 minutes. The supernatant is discarded and the inside of the tube wiped with a Kimwipe. The mitochondria is washed by re-suspending the pellet in 1X MS and repeating the 17,000 g sedimentation. The supernatant is discarded and the pellet is resuspended in a buffer. Mitochondria can be stored at -80°C for prolonged periods, e.g., up to a year, but preferably will be used shortly thereafter for NT.

This basic protocol can be modified. In particular, it may be desirable to further isolate mitochondrial DNA and use same for NT. In such case, contamination with nuclei, not small organelles, potentially is a problem and the following modifications may be made. For example, the cells may be harvested in stationary growth phase when the fewest cells are actively dividing, and CaCl₂ substituted for MgCl₂ in the RSB to stabilize the nuclear membrane. The washing of the mitochondrial pellet is omitted as is the density gradient purification. Instead, the mitochondrial pellet is simply resuspended and lysed, and the mitochondrial DNA purified from any remaining nuclear DNA. As noted, suitable methods for purifying mitochondria and mitochondrial DNA are well known in the art.

Homogenization works best if the cells are resuspended in at least 5-10X the volume of the cell pellet and if the cell suspension fills the homogenizer at least half full. Press the homogenizer pestle straight down the tube, maintaining a firm, steady pressure.

The Dounce homogenizer disrupts swollen tissue culture cells by pressure change. As the pestle is pressed down, pressure around the cell increases; when the cell slips past the end of the pestle, the sudden decrease in pressure causes the cell to rupture. If the pestle is very tight fitting, there may be some mechanical breakage as well.

B. Isolation of Mitochondria From Tissue

A mitochondrial isolation protocol is selected based on the particular tissue. For example, the homogenization buffer should be optimized for the tissue, and the optimal way to homogenize the tissue utilized. Suitable methods are well known in the art.

Rat liver is the most frequently used tissue for mitochondrial preparations because it is readily available, is easy to homogenize, and the cells contain a large number of mitochondria (1000-2000 per cell). For example, a motor-driven, Teflon and glass Potter-Elvehjem homogenizer can be used homogenize rat liver. Alternatively, if the tissue is soft enough, a Dounce homogenizer with a loose pestle can be used. The yield and purity of the mitochondrial preparation is influenced by the method of preparation, speed of preparation, and the age and physiological condition of the animal. As noted, methods of purifying mitochondria are well known.

Preferably, the buffer, tubes, and homogenizer will be pre-chilled. Pre-chilling a glass and Teflon type homogenizer creates the proper gap between the tube and pestle. The centrifugation steps are preferably effected at 40°C.

Essentially, the process will comprise removal of the liver, taking care not to rupture the gall bladder. This is placed in a beaker on ice and any connective tissue is

removed. The tissue is recognized and returned to the beaker, e.g., using very sharp scissors, a scalpel, or razor blade, mince it into 1-2 slices. The pieces are then rinsed, preferably twice, with homogenization buffer (1X MS) to remove most of the blood, and the tissue transferred to the homogenizer tube. Enough homogenization buffer is added to prepare a 1:10 (w/v) homogenate.

Use of Isolated Mitochondria or Mitochondrial DNA to Enhance NT Efficacy

It is theorized by the inventors that the efficacy of cross-species nuclear transfer may be enhanced by introduction of mitochondria or mitochondrial DNA at the same species as donor cell or nucleus. Thereupon, the nucleus DNA of resultant NT units will be species compatible.

Mitochondria isolated by the above or other known procedures are incorporated, typically by injection, into any of the following (in the case of human donor cell/bovine oocyte nuclear transfer):

- (i) non-activated, non-enucleated bovine oocytes;
- (ii) non-activated, enucleated bovine oocytes;
- (iii) activated, enucleated bovine oocytes;
- (iv) non-activated, fused (with human donor cell or nucleus) bovine oocytes;
- (v) activated, fused and cleaved reconstructed (cow oocyte/human cell) embryo; or
- (vi) activated, fused one cell reconstructed (cow oocyte/human cell) embryo.

The same procedures will enhance other cross-species NTs. Essentially, mitochondria will again be introduced into any of (i)-(vi) of the same species as the donor cell or nucleus, and the oocyte will be of a different species origin. Generally about 1 to 200 picoliters of mitochondrial suspension are injected into any of the above. The introduction of such mitochondria will result in NT units wherein the mitochondrial and donor DNA are compatible.

EXAMPLE 3

Another method for improving the efficacy of the cross-species nuclear transfer comprises the fusion of one or more enucleated somatic cells, typically human (of the same species as donor cell or nucleus), with any of the following:

- (i) non-activated, non-enucleated (e.g., bovine) oocyte;
- (ii) non-activated, enucleated (e.g., bovine) oocyte;
- (iii) activated, enucleated (e.g., bovine) oocyte;
- (iv) non-activated, fused (with human cell) oocyte (typically bovine);
- (v) activated, fused and cleaved reconstructed (e.g., cow oocyte/human cell) embryo;
- (vi) activated, fused one cell reconstructed (cow oocyte/human cell) embryo;
- or
- (vii) non-activated, fused (e.g., with human cell) oocyte (typically bovine oocyte).

Fusion is preferably effected by electrical pulse or by use of Sendai virus. Methods for producing enucleated cells (e.g., human cells) are known in the art. A preferred protocol is set forth below.

Enucleation Procedures:

5 Methods for the large-scale enucleation of cells with cytochalasin B are well known in the art. Enucleation is preferably effected using the monolayer technique. This method uses small numbers of cells attached to the growth surface of a culture disc and is ideal if limited numbers of donor cells are available. Another suitable procedure, the gradient technique, requires centrifugation of cells through Ficoll gradients and is best suited for enucleation of large number ($>10^7$) of cells.

10 ***Monolayer Technique.*** The monolayer technique is ideal for virtually any cell which grows attached to the growth surface.

15 Polycarbonate or polypropylene 250-ml wide-mouth centrifuge bottles with screw-top caps are sterilized by autoclaving. The caps preferably are autoclaved separately from the bottle to prevent damage to the centrifuge bottle. The bottle are prepared for the enucleation procedure by the sterile addition of 30 ml DMEM, 2 ml bovine serum, and 0.32 ml cytochalasin B (1 mg/ml) to each. The caps are placed on the bottles, and the bottles are maintained at 37° prior to use.

20 The cells to be enucleated (from a few hundred to $\sim 10^5$ cells) are seeded on a culture dish (35 x 15 mm; Nunc Inc., Naperville, IL). Typically, the cells are grown for at least twenty-four hours on the dishes to promote maximal attachment to the growth

surface. Preferably, the cells are prevented from becoming confluent. The culture dish is prepared for centrifugation by wiping the outside of the bottom half of the dish (containing the cells) with 70% (v/v) ethanol for the purpose of sterilization. Alternatively, the dish can be kept sterile during cell culturing by maintaining it within a larger, sterile culture dish. The medium is removed from the dish and the dish (without top) is placed upside down within the centrifuge bottle.

The rotor (GSA, DuPont, Wilmington, DE) and centrifuge are preferably pre-warmed to 37° by centrifugation for 30-45 minutes at 8000 rpm. The HS-4 swinging-bucket rotor (DuPont) can alternatively be used. The optimal time and speed of centrifugation varies for each cell type. For myoblasts and fibroblasts, the centrifuge bottle with the culture dish is placed in the pre-warmed rotor and centrifuged for approximately 20 minutes (interval between the time when the rotor reaches the desired speed and the time when the centrifuge is turned off). Preferably, speeds of 6500 to 7200 rpm are used.

After centrifugation, the centrifuge bottle is removed from the rotor, and the culture plate is removed from the bottles with forceps. A small amount of medium is maintained in the plate to keep the cells moist in order to maintain cell viability. The outside of the dish, including the top edge, is wiped with a sterile wiper, then moistened with 95% (v/v) ethanol, to remove any medium and to dry it. A sterile top is placed onto the dish. If the enucleated cells are not going to be used immediately, complete culture medium (medium supplemented with the appropriate concentration of serum) should be

While the present invention has been described and illustrated herein by reference to various specific materials, procedures, and examples, it is understood that the invention is not restricted to the particular material, combinations of materials, and procedures selected for that purpose. Numerous variations of such details can be implied and will be appreciated by those skilled in the art.

5